

NOAA National Centers for Coastal Ocean Science Center for Coastal Monitoring and Assessment

Kimani Kimbrough
W. Edward Johnson
Annie Jacob
Michael Edwards
Erik Davenport
Gunnar Lauenstein
Thomas Nalepa
Mike Fulton
Anthony Pait



February 2014



Citation for this Report

K. Kimbrough, W. E. Johnson, A. Jacob, M. Edwards, E. Davenport, G. Lauenstein, T. Nalepa, M. Fulton and A. Pait. Mussel Watch Great Lakes Contaminant Monitoring and Assessment: Phase 1. Silver Spring, MD. NOAA Technical Memorandum NOS NCCOS 180, 113 pp.



recommendation for their use by the United States government.

Mussel Watch Great Lakes Contaminant Monitoring and Assessment: Phase 1

Acknowledgements

This report could not have been completed without the cooperation, time and effort contributed by many, whose collective input has resulted in a document far superior to that which we envisioned on our own.

Dennis Apeti National Oceanic & Atmospheric Administration
Paula Bizot National Oceanic & Atmospheric Administration

Joseph Bohr Water Resources Division, Michigan Department of Environmental Quality

Marsha Burzynski Wisconsin Department of Natural Resources
Clifford Cosgrove National Oceanic & Atmospheric Administration
Dennis Donahue Great Lakes Environmental Research Laboratory,
National Oceanic & Atmospheric Administration

Andrew Fayram Wisconsin Department of Natural Resources

Mark Flipski New York State Department of Environmental Conservation

Rebecca Held
Tom Joyce

National Oceanic & Atmospheric Administration
Great Lakes Environmental Research Laboratory,
National Oceanic & Atmospheric Administration

David Klarer Ohio Department of Natural Resources

James Lazorchak Office of Research and Development, US Environmental Protection Agency

Andrew Mason National Oceanic & Atmospheric Administration
Kevin Mcmahon National Oceanic & Atmospheric Administration

Elizabeth Murphy Great Lakes National Program Office, US Environmental Protection Agency

Greg Piniak National Oceanic & Atmospheric Administration

Joseph Rathbun Water Resources Division, Michigan Department of Environmental Quality

Michael Taetsch Great Lakes Environmental Research Laboratory,


National Oceanic & Atmospheric Administration

Scott Winkler Ohio Environmental Protection Agency



Table of Contents

| 1. Executive Summary | I |
|-------------------------------------|-----|
| 2. Project Overview | 1 |
| 3. Methods | 6 |
| 4. Results | 10 |
| Analyte Characterization | 16 |
| Trace Elements Organic Contaminants | |
| Site Characterization | 52 |
| U.S. Areas of Concern Sites | 90 |
| 5. References | 112 |





Executive Summary

Recovery and rehabilitation of the Great Lakes requires the availability of sound water quality data to formulate effective, science-based policies and develop appropriate management actions. The International Joint Commission has identified monitoring as a critical component in the management and protection of the nearshore zone of the Great Lakes. The NOAA National Centers for Coastal Ocean Science (NCCOS) Mussel Watch Program (MWP) is one of the few active biomonitoring programs with a basin-wide presence and a record of consistent monitoring in the nearshore zones for close to two decades. In keeping with the mission of NCCOS to support healthy and resilient coastal communities, MWP initiated monitoring the status and trends of chemical contaminants in the Great Lakes in 1992, using dreissenid mussels as sentinel organisms. Mussels are abundant in the outer harbors on breakwaters and other hard substrates throughout the Great Lakes. These stationary, filter-feeding bottom dwellers accumulate contaminants, possess limited ability to metabolize them, are representative of local conditions, and shed light on availability of contaminants to move up the food chain, thus making them an excellent tool for contamination monitoring and assessment.

Following the inception of the Great Lakes Restoration Initiative (GLRI) to address the significant environmental issues plaguing the Great Lakes region, MWP expanded its regional monitoring activities in the Great Lakes in 2009 to include all U.S. Areas of Concern (AOCs). This enhanced effort falls under the "Toxic Substances and Areas of Concern" focus area outlined in the GLRI Action Plan. The overall objective of the expanded monitoring is to assess remediation effectiveness, and provide relevant biological data to support decision making for "Beneficial Use Impairment" (BUI) removal and subsequent delisting of AOCs. The stated objective is being addressed through a phased approach utilizing a suite of mussel and sediment indicators at newly established sites within AOCs, and by leveraging the long-term monitoring data at reference sites for meaningful interpretation of AOC status and recovery.

Herein, we report the results from the first phase under GLRI, mainly mussel tissue and sediment chemistry data obtained by MWP, with the goal of providing a high-level basin-wide perspective on the relative magnitude and extent of contamination in AOCs. This was achieved by comparing:

- 1) tissue chemistry from AOCs to reference sites;
- 2) sediment chemistry from AOCs to both MWP Great Lakes reference sites and coastal marine sites distributed nationwide and;
- 3) sediment chemistry to relevant thresholds.

Due to the availability of dreissenid mussels on and near breakwaters, this baseline assessment focused on river mouths of AOCs, and provides the foundation upon which subsequent intensive (spatially and temporally) AOC assessments will be carried out. Through numerous partnerships, MWP has begun to incorporate innovative techniques and methods in select AOCs. The results from these intensive sampling efforts will be the focus of subsequent reports. This first report sheds light on the bioavailability of select contaminants from a proven bioindicator and demonstrates the value of long-term monitoring programs in making informed coastal ecosystem management decisions.

Thank you,

The Great Lakes Mussel Watch Team



National Centers for Coastal Ocean Science

Project Overview

- With Great Lakes Restoration Initiative funds, NOAA Mussel Watch expanded monitoring activities into all U.S. Areas of Concern (AOCs) and added additional analyses to address Beneficial Use Impairment (BUI) target goals and effectiveness of AOC restoration efforts.
- All U.S. AOCs were sampled for mussel tissue and/or sediment as part of the Phase 1 Great Lakes baseline study.
- Historic Mussel Watch sites, primarily located in shallow (<20 ft) nearshore lake water, were used as reference sites for comparison with AOCs.
- National sediment sites were used to bring perspective to Great Lakes sediment concentrations.
- Mussel contaminant concentrations provided an assessment of bioavailable contamination inside and outside AOC boundaries.

The Great Lakes constitute the largest freshwater system in the world, containing more than 90% of the nation's, and 20% of the world's freshwater supply. The Great Lakes basin is home to over 35 million people and supports numerous economic activities including agriculture, industries, commercial and sport fisheries, tourism, and recreation. Exploitation of the Great Lakes has resulted in multiple environmental challenges and has negatively impacted many beneficial uses of this globally treasured natural resource.

In 2009, the Great Lakes Restoration Initiative (GLRI) was signed into law to restore and rehabilitate the designated areas in the Great Lakes. This initiative is guided by an action plan created by a task force of 11 federal agencies (GLRI Action Plan, 2010) and identifies five focus areas:

- 1. Toxic substances and Areas of Concern;
- 2. Invasive species;
- 3. Nearshore health and nonpoint source pollution;
- 4. Habitat and wildlife protection and restoration; and
- 5. Accountability, education, monitoring, evaluation, communication and partnerships.

While considerable progress has been made in improving the condition of the Great Lakes through environmental legislation and the Great Lakes Water Quality Agreement between the United States and Canada (GLWQA, 2012), pollutant loadings from point and non-point sources, and legacy contaminants in sediment continue to impair the Great Lakes ecosystem. This report focuses on the legacy contamination in U.S. AOCs, which are defined in the GLWQA as "a geographic area designated by the Parties where significant impairment of beneficial uses has occurred as a results of human activities at the local level." There are 31 AOCs on the U.S. side (5 of those shared with Canada) with one or more Beneficial Use Impairments (Table 1; Figure 1). BUIs have to be removed for the AOCs to be delisted. AOC specific

BUIs are too numerous to list here but can be found online at http://www.epa.gov/glnpo/aoc/.

The National Oceanic and Atmospheric Administration (NOAA) Mussel Watch Program (MWP) administered by the National Centers for Coastal Ocean Science (NCCOS) has used dreissenid mussels to monitor contaminants in the nearshore zones of the Great Lakes since 1992. This contaminant monitoring effort in the Great Lakes is part of a larger contaminant monitoring program that began in 1986 and currently includes more than 300 sites nationwide (Figure 2). In the Great Lakes, MWP established 7 sites in the inaugural year and more sites were added in the initial years after the invasion and proliferation of Ponto-Caspian mussels *Dreissena polymorpha* and *Dreissena* rostriformis bugensis (zebra and quagga mussels, respectively) that became established in the region in the late 1980s (Carlton 2008; Mills et al., 1993; Robertson and Lauenstein, 1998). The abundance and widespread distribution of dreissenid mussels in the nearshore areas of the Great Lakes (except Lake Superior) renders them ideal candidates for routine contaminant monitoring.

Dreissenid mussels are widely used as bioindicators in Europe and their native Ponto-Caspian region, because of their high filtering rates, high lipid content and the ability to bioaccumulate contaminants from water, algae and suspended sediments (de Kock and Bowmer 1993; Bervoets et al., 2005; Berny et al., 2003; Gossiaux et al., 1998; Reeders and bij de Vaate 1992). Other favorable characteristics of dressenids as biomonitors include ease of collection, abundance, sessile life cycle, limited ability to metabolize lipophilic contaminants, and utility in effects-based monitoring. In addition, the recognition of dressenid mussels as keystone species of the Great Lakes ecosystem (Vanderploeg et al., 2002), reinforces its critical value in the assessment of the overall health and ecological forecasting of the Great Lakes.

Table 1. There were 31 U.S. Areas of Concern (AOCs) in the U.S. Great Lakes suffering 1 or more of 14 Beneficial Use Impairments (BUIs) listed. MWP provides data and information to potentially support the decision making to remove select BUIs.

| Restrictions on fish & wildlife consumption | Eutrophication or undesirable algae |
|---|--|
| Tainting of fish and wildlife flavor | Restrictions on drinking water |
| Degradation of fish & wildlife populations | Beach closings |
| Fish tumors or other deformities | Degradation of aesthetics |
| Bird or animal deformities or reproduction problems | Added costs to agriculture or industry |
| Degradation of benthos | Degradation of phytoplankton & zooplankton populations |
| Restrictions on dredging activities | Loss of fish & wildlife habitat |

Project Approach for Phase 1: Basin-Wide Baseline Assessment

- Established sites in all U.S. AOCs, and assessed contaminants in mussels and sediments in these new sites;
- Leveraged data, information, and knowledge from long-term Mussel Watch sites in Great Lakes (1992-present) and nation (1986-2009) and evaluated the extent and magnitude of pollution in AOCs relative to long-term Mussel Watch monitoring sites (i.e. reference sites).
- Determined baseline chemical contaminant levels at AOCs and reference sites, to compare regional, lake-wide, and local progress toward AOC chemical reduction targets.

Currently, there are 25 long-term Mussel Watch sites in the Great Lakes from Duluth, MN on Lake Superior at the mouth of the St. Louis River to Cape Vincent, NY where Lake Ontario flows into the St. Lawrence River. The majority of these sites were established away from known impacted areas, following the overall program directive to measure contamination at sites that represent ambient concentrations. These long-term monitoring sites thus served as reference sites for contamination assessment in highly impacted areas (e.g., AOCs) in the Great Lakes.

Aligning with the focus of GLRI, MWP expanded its monitoring activities in the region to include all U.S. AOCs (Figure 1). The overall objective of the expanded monitoring was to provide relevant biomonitoring data to potentially support decision making for BUI removal and subsequent delisting of U.S. AOCs. The stated objective is being addressed through a two phase approach utilizing mussel and sediment

indicators. Phase 1 effort, the basin-wide assessment of mussels and sediment at newly established sites within U.S. AOCs and long-term Mussel Watch sites (i.e. reference sites) is presented in this report. Phase 2 has two ongoing parts consisting of intensive mussel sampling in select U.S. AOCs (where remediation is imminent or has already occurred) for improved spatial coverage, and probabilistic ecosystem assessments to characterize sediment quality in select AOCs. As part of the Phase 2 intensive mussel sampling initiative, newer biomonitoring approaches including the use of caged mussels for contaminant source tracking and effects-based monitoring to assess mussel health have been incorporated.

In this report, we summarize contaminant chemistry data in mussels and sediment for the Phase 1 basin-wide assessment and leverage two decades of historic Mussel Watch Program data from the Great Lakes and the Nation (Figure 2) for comparison.




Figure 1. Mussel Watch Great Lakes baseline assessment U.S. AOC sites (labeled).




Figure 2. National Mussel Watch sites.

The basin-wide baseline data presented here can be used by stakeholders and resource managers for AOC remediation effectiveness assessment and as an additional biological line of evidence to support BUI removal. Specifically, mussel tissue concentrations could potentially be used to set target chemical reduction goals for remediation purposes as these mussels are robust indicators of biovailable contamination. Further, mussels serve as a potential additional line of evidence to support decision making with respect to "Restriction on fish and wildlife consumption" BUI. Unlike fish, which move in and out of AOC boundaries, mussels bioaccumulate and integrate contaminant exposure at the specific location where they are found in an AOC. More importantly, mussels have limited ability in metabolizing xenobiotics, including PAHs and PCBs, compared to fish (Farrington, 1983).

The role of dreissenid mussels in contaminant cycling and biomagnification of pollutants (indirectly via deposition of feces and psuedofeces, and directly via predation in the food chain) in the Great Lakes is well documented (Bruner et al., 1994a, b; Morrison et al., 1998; Marvin et al., 2000; Cho et al., 2004). Monitoring contaminants in dreissenids can thus not only reflect ambient contamination levels, but can also further our understanding of trophic transfer and biomagnification.

Recently, the International Joint Commission (IJC 2011) emphasized the need for a greater understanding of nearshore zone processes and management, subsequent offshore effects, and the health of the lakes as a whole. To these ends, monitoring of chemical, biological, and toxicological indicators in dreissenid mussels provides a means to further assess temporal trends, biological uptake, and ultimate fate of many contaminants.

Current & Future Work

Our baseline monitoring effort in 2009/2010 was in essence an expansion of the fixed sampling design of traditional Mussel Watch and provides a synoptic view of the relative magnitude and extent of contamination in the harbor areas of AOCs and reference sites. In order to better address contamination and remediation issues of specific AOCs, Mussel Watch has adopted multi-parameter assessments and incorporated newer techniques and approaches to the existing methods. For mussels, we have intensified sampling, both spatially and temporally, in select AOCs (e.g., Sheboygan River AOC; Figure 3) to provide a more robust measure of bioavailable contamination, have utilized deployment of caged mussels to source track contaminants in areas where extant mussels beds are not found (Figure 4), and have incorporated effects-based monitoring to complement body burden




Figure 3. Densification of mussel and sediment samples temporally and spatially.

measurements. For the sediment matrix, probabilistic sediment quality assessment was carried out to report on sediment chemistry, benthos and sediment toxicity, which could directly address Degradation of Benthos BUI. These focused efforts in select AOCs have been implemented through successful partnerships and collaborations with the local, state and federal entities. The results from these ongoing and future work will be the focus of subsequent reports.



Figure 4. Caged mussels, passive samplers and Hester Dendys being retrieved after deployment for 6 weeks in Manistique River AOC.



U.S. Great Lakes Areas of Concern Assessment



Mussel Watch sampled sites in 31 U.S. AOCs as part of the basin-wide baseline assessment under GLRI (Figure 1) in 2009 and 2010. All mussel sites were established at river mouths or outer harbor areas where stone breakwaters provided a colonizing substrate for dreissenid mussels. When available, one mussel station and one sediment station per site were collected from each site in the eastern Great Lakes (downstream from Detroit) in 2009. In the western Great Lakes, one mussel station and up to three sediment stations were collected per site in 2010. The Grand Calumet River and St. Louis River AOC are the exception, each had 2 baseline mussel sites. One composite sample comprising 50-100 mussels was sampled from each sation. Only sediment samples were collected at Torch Lake, Carp River, St. Marys River, Kalamazoo River and Rouge River.

Contaminant monitoring of mussels and sediment in the Great Lakes by MWP began in 1992. Prior to GLRI, MWP was monitoring 25 sites routinely in the Great Lakes. Data from long-term MWP sites that were located in AOCs (Milwaukee Estuary, Niagara River, Rochester Embayment, Saginaw River and Bay, and Grand Calumet River) were included in AOC results throughout this document. Long-term MWP sites located outside of AOCs were used as reference sites and were compared to AOCs (Figure 5). The reference sites were located within watersheds that represented a variety of land uses and occured within the shallow, nearshore zones (< 20 ft).

Within a MWP site, sediment stations were co-located with mussel stations. Some sites did not have fine-grained (percent silt-clay) sediments and hence sediment sites were relocated nearby, consistent with procedures of the National Coastal Assessment Quality Assurance Project Plan 2001-2004 (US EPA 2001). Sites were sampled on a biennial basis; Lakes Ontario and Erie sites were sampled in odd years, and Lakes Huron, Superior and Michigan sites in even years.

Sample Collection

Mussels were collected from natural substrates by diving or dredging. Upon collection, mussels were rinsed with site water to remove debris, placed in freezer bags, packed on water ice and shipped to laboratories within two days. Samples for sediment chemistry, toxicity, and the benthic community analyses were collected concurrently with a Young-modified, van Veen grab sampler (0.04 m²) deployed from a small boat. Protocols for sample collection, preparation and analysis are found in the Quality Assurance Project Plan and in Lauenstein and Cantillo (1998 and references therein), Kimbrough and Lauenstein (2006), Kimbrough et al., (2006), and available online at http://coastalscience.noaa.gov/projects/detail?key=179




Figure 5. Mussel Watch reference sites (labeled).

Table 2. Contaminants summarized as part of the Expanded Great Lakes MWP. The complete list of contaminants and associated method detection limits can be downloaded at http://coastalscience.noaa.gov/projects/detail?key=179

| Compound class | Compound |
|---|---|
| Trace elements | Arsenic, cadmium, copper, lead, methyl mercury, mercury, zinc |
| Chlordane | Alpha-chlordane |
| Chlorpyrifos | Chlorpyrifos |
| Dichlorodiphenyltrichloroethane and metabolites (DDT) | 2,4'-DDD; 2,4'-DDE; 2,4'-DDT; 4,4'-DDD; 4,4'-DDE; 4,4'-DDT |
| Heptachlor | Heptachlor |
| Hexachlorobenzene | Hexachlorobenzene |
| Mirex | Mirex |
| Polycyclic aromatic hydrocarbons (PAHs) | Benzo[a]pyrene, benzo[e]pyrene |
| Polychlorinated biphenyls (PCBs) | PCB8/5, PCB18, PCB28, PCB 29, PCB31, PCB44, PCB45, PCB49, PCB52, PCB56, PCB66, PCB70, PCB74, PCB87, PCB95, PCB99, PCB101/90, PCB105, PCB110, PCB118, PCB128, PCB138, PCB146, PCB149, PCB151, PCB153/132/168, PCB156, PCB158, PCB170/190, PCB174, PCB180, PCB183, PCB187, PCB194, PCB195/208, PCB199, PCB201, PCB206, PCB209 |

Contaminants

The recently established AOC sites together with long-term MWP reference sites provide relevant data to aid in the cleanup and subsequent delisting of AOCs. All sites were monitored for a wide array of contaminants, including trace elements and legacy organic contaminants. For the purpose of this report, only select and relevant analytes are presented (Table 2); however, data for additional analytes can be downloaded (http://coastalscience.noaa.gov/projects/detail?key=179). Ancillary measurements for mussels and sediment were also recorded. For mussels, these consisted of gonadal index, wet weight, dry weight, percent lipid, and shell length and volume. For sediment, these consisted of grain size (percent sand, silt and clay) and total organic carbon.

Data Quality

The Mussel Watch Program used a performance based quality assurance process to ensure data quality. This effort has been in operation since 1985 and is designed to document sampling protocols, analytical procedures and laboratory performance. Analytical laboratories used by the Mussel Watch Program are required to participate in exercises with assistance from the

National Institute of Standards and Technology (NIST) and the National Research Council of Canada (NRC) to ensure data are comparable in accuracy and precision (Willie, 2000; Schantz et al., 2000).

Statistical Analysis

Tissue and sediment contaminant concentrations from all the AOC and reference sites were assigned to a concentration range (high, medium and low) using hierarchical Wards cluster analysis. This analysis allowed clustering of contaminant concentrations into groups such that the numbers contained within a group are more like each other than any other number in a different group. Results in the low, medium and high categories did not represent measurements that have exceeded regulatory thresholds; rather, it denoted that they are significantly higher than the preceding category (p < 0.05; Wilcoxon). The high, medium and low categories are not associated with human health or ecosystem health endpoints. In cases where the concentration measurements could not be grouped into a category of at least 3, the measurements were labeled as outliers, and removed to perform the analysis again.


The Shapiro-Wilks test was used to test for normality. None of the analytes tested had a normal distribution so non-parametric statistics were used. The nonparametric Kruskal Wallis test was used to test for significant differences in contaminant concentrations among AOCs and reference sites, and between national and Great Lakes (inclusive of AOC and reference) sites. Data summaries (descriptive statistics) for selected chemicals measured in mussel tissue and sediment are provided categorically by chemical and by site. Descriptive statistics are organized by site type (AOC and reference site), and by matrix (mussel and sediment). Sediment results for the Great Lakes are compared to the latest national MWP survey (2006/2007). Concentration of contaminants in dreissenid mussel tissue from the Great Lakes were not compared to concentrations found in mussels/ bivalves from other regions because of contaminant uptake and bioaccumulation differences among species.

Benchmarks and Background


Resource managers rely on chemical-specific, numerically based benchmarks to guide their actions to ensure protection of the environment and/or human health. This report uses the term "benchmark" to describe any published chemical concentration or concentration range reported in sediment (e.g., "criteria," "standard," guideline," value," indicator," "alert or action level," "tolerance level," "threshold level").

Sediment benchmarks commonly reported in the literature as sediment quality guidelines (SQGs) rely on empirical or theoretically based approaches that are predictive of toxicity and bioaccumulation. For this report, we used freshwater sediment quality guidelines reported by MacDonald et al., (2000), their consensusbased numbers, Probable Effects Concentration (PEC), and Threshold Effects Concentration (TEC). The PECs are SQGs for individual chemicals above which adverse effects in sediments are expected to frequently occur, whereas TECs were developed to provide an estimate of conditions where toxicity would not be expected. If no SQG was reported by MacDonald et al., (2000), NOAA SQuiRTs (Screen Quick Reference Tables) were consulted, followed by the Canadian **Environmental Quality Guidelines.**

Benchmarks for biota, including shellfish, are scarce. The FDA provides "action levels" and "tolerance levels" for fish and shellfish for a few chemicals but these are intended for regulating food safety and commerce. The EPA provides "screening levels" for management decisions to issue fish consumption advisories, but these are of limited value for assessing mussel contaminant levels. In this study, we have not attempted to compare dreissenid mussel tissue concentration to any existing thresholds for other bivalve tissue. Instead, the mussels tissue concentrations in U.S. AOCs were interpreted by leveraging data from reference sites.



U.S. Great Lakes Areas of Concern Assessment



Major results and discussion, based on statistical comparisons of AOCs, references sites, and coastal marine sites, are presented in the ensuing paragraphs. Graphical summarization of results by analyte and by site appear in the Analyte and Site Characterization sections respectively.

AOC vs. Reference Sites

Data analyses indicated that greater than 50 percent of all tissue and sediment measurements (AOC and reference sites) were categorized as low, and greater than 85 percent of all measurements were categorized as low or medium (Figure 6). Conversely, high and outlier sites together represented 5-15 percent of all sediment and tissue measurements (Figure 6). The difference between AOC and reference sites was primarily found in the percentage of each that comprise the elevated measurements found in the high and outlier categories. For trace elements and organics in sediment, and organics in tissue, a higher proportion of high and outlier measurements were found at AOC sites (Figure 6). In contrast, the percentage of outlier and high measurements were almost equal for trace elements in mussel tissue in AOC and reference sites.

For tissue, no significant difference was found between AOC and reference sites for trace elements examined except for cadmium. Cadmium concentrations in mussels from reference sites were higher than those from AOC sites (Kruskal Wallis; p <0.05; Figure 7). Four of the nine organic analytes (DDT, Benzo[a] pyrene, Benzo[e]pyrene and PCB) measured in dreissenid tissue were significantly higher at AOCs




Figure 6. Percentage of low (green), medium (yellow), high (red), and outliers (maroon) categories for metals and organics in AOCs and reference sites (Ref.) for both tissue and sediment matrices.

when compared to reference sites (Kruskal Wallis; p <0.05; Figure 7). The lack of significance among other analytes may be due to the fact that majority of sites (both AOC and reference sites) have concentrations that fall in the low and medium categories.




Figure 7. Comparison of AOC and reference sites tissue measurement means with standard error bars. Asterisk represent analytes with significant differences. Organic compounds are lipid normalized.




Figure 8. Comparison of AOC and reference sites concentration means with associated standard error bars.

For sediment, there were no significant differences found between AOCs and reference sites for organics or trace elements (Kruskal Wallis; p > 0.05; Figure 8).

Great Lakes vs. National Sediment Sites

The sediment concentrations for all analytes from Great Lakes were higher than the concentrations from the rest of the MWP sites in the nation (Kruskal Wallis; p <0.05; Figure 9). All of the trace elements and organic contaminants examined were higher in the Great Lakes with respect to the nation as a whole irrespective of

inclusion of AOCs in the analysis (Figure 9), that is, even Great Lakes reference sites were higher.

MWP's national sediment measurements are obtained from small embayments, estuaries, and open coast sites, many of which have higher flushing/turnover rates when compared to the Great Lakes. Conversely, the residence time of some of the persistent pollutants in the Great Lakes is higher, which may partly explain the difference between national and Great Lakes sediment measurements.




Figure 9. Comparison of Great Lakes (AOCs and reference sites) and national sediment concentration means with associated standard error bars. Asterisk represent analytes with significant differences.

Comparison Among AOCs

To identify the relative magnitude and extent of contamination in the outer harbor areas among AOCs, heat maps were generated for both tissue and sediment, which show the heirarchical Ward's cluster analysis categories (outlier, high, medium and low). These heat maps summarize the data in maps and associated figures in Section 2 & 3, and clearly show the utility of using one unique matrix for characterizing sites as well as dentifying sites with elevated concentrations that require further investigation. However, unlike abiotic measurements. mussels integrate the contaminant signal over long periods of time allowing for a time-integrated indication of environmental contamination. More importantly, contaminant tissue burden in mussels that reside at the base of the food chain reflect contaminant bioavailability and bioaccumulation potential, and thus provide information about environmental contamination and effects that cannot be defined by abiotic matrices. Therefore, having mussel tissue data from AOC sites across the basin provide biologically relevant data that can be used in making informed management decisions.

Readers should be mindful that these results are not representative of the entire AOC and that low concentrations in the harbor area (where the majority of the sites are located) do not imply that the entire AOC is free of contamination. However, this baseline assessment effort helps identify the contaminant concentration in the harbor, which can serve as a chemical reduction target goal for remediation purposes. To provide enhanced characterization of AOCs, intensive sampling is required, and has begun at some priority AOCs.

Benchmarks

Specific benchmarks and threshold concentrations were not found for dreissenid mussels. As such, mussel tissue concentrations from AOCs were compared to concentrations obtained from reference sites. In contrast, for sediment, consensus based TEC and PEC sediment quality guidelines (MacDonald et al., 2000) was used to compare concentrations from AOCs and reference sites. For several chemicals (DDT, PCBs, benzo(a)pyrene), the mean reference site concentration exceeded or was only slightly below the


reported TEC. Other chemicals with TEC exceedances at many sites included arsenic, cadmium, copper, lead, total mercury, zinc, chlordane (as alpha chlordane), DDT, and heptachlor epoxide (as heptachlor).

Interpretation of Chemical Data


Trace elements are naturally occurring but can have anthropogenic sources as well. Distinguishing the sources of elevated concentrations is beyond the scope of MWP, however, our monitoring data can identify sites with elevated concentrations where further investigation is required.

The organic compound results have two primary patterns of distributions. The first distribution is characterized by the majority of the sites being at or near detection limits with a few sites exhibiting measurements that are elevated, (chlordane, chlorpyrifos, heptachlor, hexachlorobenzene, Mirex); essentially a presence and absence distribution. The second group of organic compounds are more ubiquitous and thus have a "background level" arising from its chemical persistence and also have several outlier measurements that are orders of magnitude higher than the "background" measurements (Benzo [a]pyrene, Benzo[e]pyrene, DDT, and PCB). This second group of organic contaminants approaches a skewed distribution that is similar to the trace element distribution (Site Characterization Section).

Most regulatory controls of chlorinated organics went into effect in the 1970s and 1980s, prior to the initiation of mussel monitoring in the Great Lakes. The tissue trend map for reference sites (Analyte Characterization Section-Page 16) shows historic site measurements for a subset of MWP long-term data. The box and whisker plots for organic contaminants from reference sites show no significant difference between years since 2001 across the entire basin, and is supported by the bar charts in the adjacent map. These results suggest that decreasing organic contaminant trends have reached asymptotic levels and further reductions in these levels by natural means alone will likely take decades.



Tissue heat map. Figure showing outlier (★), high (●), medium (○), and low (●) AOC tissue measurements for all trace elements, and organic contaminants discussed in this report.



Sediment heat map. Figure showing outlier (★), high (●), medium (○), and low (○) AOC sediment measurements for all trace elements and organic contaminants discussed in this report. Boxes are white when no data is available.




Analyte Characterization

- Great Lakes sediment (U.S. AOC & reference) measurements were found to be significantly higher than national measurements.
- Significant differences between U.S. AOC and reference site tissue measurements were primarily limited to organic contaminants.
- When compared to national and reference site measurements, U.S. AOCs registered
 a disproportionate number of outlier and high measurements for organic contaminants
 but not for trace elements.
- Sediment measurements for U.S. AOCs were not significantly different from reference sites sediment measurements.







U.S. Great Lakes Areas of Concern Assessment




Tissue map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (μ g/g dry wt.) in dressenid mussels.




AOC barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (μ g/g dry wt.). Reference lines represent mean AOC (solid line) and reference site (dotted line) concentrations.




Historic data: Mussel tissue concentration (μg/g dry wt.) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and U.S. AOC sites collected in 2009/2010.




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (μ g/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (μ g/g dry wt.). Where available, reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map: Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; μ g/g dry wt.).




Tissue map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (μ g/g dry wt.) in dressenid mussels.




AOC barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (μ g/g dry wt.). Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data: Mussel tissue concentration (μ g/g dry wt.) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (μ g/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (μ g/g dry wt.). Where relevant, reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map: Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; μg/g dry wt.).




Tissue map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (μ g/g dry wt.) in dressenid mussels.




AOC barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (μ g/g dry wt.). Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data: Mussel tissue concentration (μ g/g dry wt.) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.



Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (μ g/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (μ g/g dry wt.). Where relevant, reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.







National frequency plot and sediment map: Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; μg/g dry wt.).




Tissue map: Categories low (\bullet) , medium (\bullet) and high (\bullet) , and where applicable outlier (\star) , are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (ng/g | lipid) in dressenid mussels.




AOC Barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (ng/g lipid) among AOCs. Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data. Mussel tissue concentration (ng/g lipid) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (ng/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (ng/g dry wt.). Where relevant reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map. Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; ng/g dry wt.). Extreme outliers omitted.




Tissue map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (μ g/g dry wt.) in dressenid mussels.




AOC barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (μ g/g dry wt.). Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data: Mussel tissue concentration (μg/g dry wt.) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (μ g/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (μ g/g dry wt.). Where relevant, reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map: Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; μg/g dry wt.).




Tissue map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (μ g/g dry wt.) in dressenid mussels.




AOC barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (µg/g dry wt.). Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (μ g/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (μ g/g dry wt.). Where relevant, reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




Tissue map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (μ g/g dry wt.) in dressenid mussels.




AOC barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (µg/g dry wt.). Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data: Mussel tissue concentration (μ g/g dry wt.) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (μ g/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (μ g/g dry wt.). Where relevant, reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map: Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; μg/g dry wt.).




Tissue map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (μ g/g dry wt.) in dressenid mussels.




AOC barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (µg/g dry wt.). Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data: Mussel tissue concentration (μ g/g dry wt.) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (μ g/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (μ g/g dry wt.). Where relevant, reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.







National frequency plot and sediment map: Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; μg/g dry wt.).




Tissue map: Categories low (\bullet) , medium (\bullet) and high (\bullet) , and where applicable outlier (\star) , are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (ng/g lipid) in dressenid mussels.




AOC Barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (ng/g lipid) among AOCs. Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data. Mussel tissue concentration (ng/g lipid) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (ng/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (ng/g dry wt.). Where relevant reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map. Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; ng/g dry wt.).




Tissue map: Categories low (\bullet) , medium (\bullet) and high (\bullet) , and where applicable outlier (\star) , are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (ng/g lipid) in dressenid mussels.




AOC Barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (ng/g lipid) among AOCs. Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data. Mussel tissue concentration (ng/g lipid) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (ng/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (ng/g dry wt.). Where relevant reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map. Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; ng/g dry wt.).




Tissue map: Categories low (●), medium (●) and high (●), and where applicable outlier (★), are used to characterize U.S. AOC (●) and reference (■) site contaminant concentrations (ng/g lipid) in dressenid mussels.




AOC Barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (ng/g lipid) among AOCs. Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data. Mussel tissue concentration (ng/g lipid) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (ng/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (ng/g dry wt.). Where relevant reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map. Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; ng/g dry wt.).




Tissue map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (ng/g lipid) in dressenid mussels.




AOC Barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (ng/g lipid) among AOCs. Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data. Mussel tissue concentration (ng/g lipid) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.



Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (ng/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (ng/g dry wt.). Where relevant reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map. Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; ng/g dry wt.). Extreme outliers omitted.


Hexachlorobenzene



Tissue map: Categories low (\bullet) , medium (\bullet) and high (\bullet) , and where applicable outlier (\star) , are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (ng/g lipid) in dressenid mussels.




AOC Barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (ng/g lipid) among AOCs. Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data. Mussel tissue concentration (ng/g lipid) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.


Hexachlorobenzene




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (ng/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (ng/g dry wt.). Where relevant reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map. Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; ng/g dry wt.).




Tissue map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (ng/g lipid) in dressenid mussels.




AOC Barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (ng/g lipid) among AOCs. Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data. Mussel tissue concentration (ng/g lipid) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.



Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (ng/g dry wt.).



AOC barchart: Comparison of AOC contaminant concentrations in sediment (ng/g dry wt.). Where relevant reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map. Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; ng/g dry wt.).

Benzo(a) pyrene


Tissue 2009/2010

1067 - 1620
38041


Tissue map: Categories low (\bullet) , medium (\bullet) and high (\bullet) , and where applicable outlier (\star) , are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (ng/g lipid) in dressenid mussels.




AOC Barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (ng/g lipid) among AOCs. Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data. Mussel tissue concentration (ng/g lipid) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (ng/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (ng/g dry wt.). Where relevant reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map. Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; ng/g dry wt.). Extreme outliers omitted.




Tissue map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (ng/g lipid) in dressenid mussels.




AOC Barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (ng/g lipid) among AOCs. Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data. Mussel tissue concentration (ng/g lipid) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.




Sediment map: Categories low (●), medium (●) and high (●), and where applicable outlier (★), are used to characterize U.S. AOC (●) and reference (■) site sediment concentrations (ng/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (ng/g dry wt.). Where relevant reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.




National frequency plot and sediment map. Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; ng/g dry wt.). Extreme outliers omitted.




Tissue map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site contaminant concentrations (ng/g lipid) in dressenid mussels.




AOC Barchart: Comparison of AOC contaminant concentrations in dreissenid mussels (ng/g lipid) among AOCs. Reference lines represent mean AOC (solid line) and reference site (dashed line) concentrations.




Historic data. Mussel tissue concentration (ng/g lipid) from reference sites collected from 2001- 2008 (map), and whisker plot of the mapped data in addition to reference and AOC sites collected in 2009/2010.




Sediment map: Categories low (\bullet), medium (\bullet) and high (\bullet), and where applicable outlier (\star), are used to characterize U.S. AOC (\bullet) and reference (\blacksquare) site sediment concentrations (ng/g dry wt.).




AOC barchart: Comparison of AOC contaminant concentrations in sediment (ng/g dry wt.). Where relevant reference lines representing PEC (black dashed line), TEC (blue dashed line), and reference site mean (gray solid line) are given.



National frequency plot and sediment map. Frequency plot of national (2006/2007) and Great Lakes (AOC and reference; 2009/2010) sediment data, and map of national sediment concentrations (2006/2007; ng/g dry wt.). Extreme outliers omitted.




Site Example




^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

U.S. Areas of Concern





St. Louis River




| Lege | nd | | | |
|------|----|----------|--------|-----------|
| Low | | Medium _ | High 🛑 | Outlier ★ |

| | | | | | | | | 1 | 34 |
|----------------|----------|------|-------------|-------|------------|----|----------|------|----|
| Metals | Base | line | - 2009/10** | Sedim | ent - 2010 | | Tissue - | 2011 | |
| (μg/g dry wt.) | Tissue 🔾 | | Sediment | | | O, | 0 | 0 | |
| Arsenic | 3.3 | | 4.5 | 5.9 | 6.7 | | | | |
| Cadmium | 1.21 | | 0.29 | 0.55 | 0.52 | | | | |
| Copper | 14 | | 21 | 30 | 34 | | | | |
| Lead | 1.7 | | 19 | 27 | 32 | | | | |
| Methyl Mercury | 0.08 | | 0.000 | 0.000 | 0.000 | | | | |
| Mercury | 0.12 | | 0.090 | 0.176 | 0.210 | | | | |
| Zinc | 120 | | 87 | 141 | 165 | | | | |

| Organics | Basel | ine - | - 2009/10** | | Sedime | nt - 2010 | | Tissue | - 2011 |
|-------------------|----------|-------|-------------|---|--------|-----------|----|--------|--------|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment | l | | | Ο, | 0 | 0 |
| Alpha-Chlordane | 8.2 | | 0.05 | | 0.25 | 0.36 | | | |
| Chlorpyrifos | 0.00 | | 0.05 | | 0.51 | 0.29 | | | |
| DDT | 130 | | 0.8 | | 3.4 | 4.7 | | | |
| Heptachlor | 0 | | 0.12 | | 0.10 | 0.06 | | | |
| Hexachlorobenzene | 4 | | 0.09 | | 0.08 | 0.14 | | | |
| Mirex | 1.4 | | 0.045 | | 0.038 | 0.024 | | | |
| Benzo[a]pyrene | 1141 | | 208 | | 741 | 435 | | | |
| Benzo[e]pyrene | 1276 | | 152 | | 446 | 287 | | | |
| PCB | 894 | | 21 | | 19 | 29 | | | |

Minnesota Point



Legend Low ● Medium ● High ● Outlier ★


| | Z / |
|--|-----|

| Arsenic | 3.9 | |
|----------------|------|---|
| Cadmium | 5.88 | • |
| Copper | 15 | |
| Lead | 0.3 | |
| Methyl Mercury | 0.02 | |
| Mercury | 0.05 | |
| Zinc | 97 | |

| Alpha-Chlordane | 5.9 | |
|-------------------|-----|---|
| Chlorpyrifos | 12 | • |
| DDT | 33 | • |
| Heptachlor | 0 | |
| Hexachlorobenzene | 11 | • |
| Mirex | 0.0 | |
| Benzo[a]pyrene | 185 | • |
| Benzo[e]pyrene | 57 | |
| PCB | 305 | |


^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Torch Lake



| Organics | Baseline | e - 2009/10** | Sedimer | nt - 2010 | | Tissue | - 2011 |
|-------------------|----------|------------------|---------|-----------|----|--------|--------|
| (ng/g dry wt.)* | Tissue 🔵 | Sediment | | | O, | 0 | 0 |
| Alpha-Chlordane | | 0.08 | 0.02 | 0.32 | | | |
| Chlorpyrifos | | 0.00 | 0.03 | 0.11 | | | |
| DDT | | 2.4 | 0.3 | 5.2 | | | |
| Heptachlor | | 0.04 | 0.03 | 0.27 | | | |
| Hexachlorobenzene | | 0.15 | 0.55 | 0.68 | | | |
| Mirex | | 0.071 | 0.000 | 0.069 | | | |
| Benzo[a]pyrene | | 42 | 7 | 173 | | | |
| Benzo[e]pyrene | | 39 | 7 | 122 | | | |
| PCB | | 25 | 3 | 43 | | | |

Carp River




| Arsenic | 6.8 | 12.4 |
|----------------|-------|-------|
| Cadmium | 0.29 | 0.00 |
| Copper | 17 | 10 |
| Lead | 21 | 11 |
| Methyl Mercury | 0.000 | 0.000 |
| Mercury | 0.002 | 0.049 |
| Zinc | 58 | 40 |

| Alpha-Chlordane | 0.00 | 0.00 |
|-------------------|-------|-------|
| Chlorpyrifos | 0.00 | 0.00 |
| DDT | 0.0 | 0.2 |
| Heptachlor | 0.00 | 0.00 |
| Hexachlorobenzene | 0.00 | 0.04 |
| Mirex | 0.000 | 0.017 |
| Benzo[a]pyrene | 1 | 66 |
| Benzo[e]pyrene | 1 | 39 |
| PCB | 0 | 0 |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

St. Marys River



| Organics | Baselir | ne - 2009/10** | Sedime | ent - 2010 | |
|-------------------|----------|----------------|--------|------------|--|
| (ng/g dry wt.)* | Tissue 🔵 | Sediment | | | |
| Alpha-Chlordane | | 0.00 | | | |
| Chlorpyrifos | | 0.00 | | | |
| DDT | | 0.0 | | | |
| Heptachlor | | 0.00 | | | |
| Hexachlorobenzene | | 0.01 | | | |
| Mirex | | 0.000 | | | |
| Benzo[a]pyrene | | 29 | | | |
| Benzo[e]pyrene | | 19 | | | |
| PCB | | 1 | | | |

Manistique River



| Lege | nd | | | |
|------|----|----------|--------|-----------|
| Low | | Medium 🛑 | High 🛑 | Outlier ★ |


| - Mary | | | |
|--------|---|--|----|
| { ** | 0 | The state of the s | |
| 3 | | 2 | |
| | | | \$ |

| Arsenic | 5.6 | | 1.7 | 0.7 | 1.1 | 7.5 | 7.6 |
|----------------|------|---|-------|-------|-------|-------|-------|
| Cadmium | 0.70 | 0 | 0.00 | 0.00 | 0.00 | 1.70 | 1.72 |
| Copper | 6 | | 4 | 2 | 3 | 26 | 108 |
| Lead | 0.3 | 0 | 12 | 6 | 10 | 3 | 4 |
| Methyl Mercury | 0.03 | | 0.000 | 0.000 | 0.000 | | |
| Mercury | 0.04 | | 0.020 | 0.009 | 0.012 | 0.073 | 0.077 |
| Zinc | 49 | | 24 | 9 | 20 | 70 | 80 |

| Alpha-Chlordane | 5.9 | | 0.00 | 0.00 | 0.00 | 6.3 | 8.7 |
|-------------------|------|---|-------|-------|-------|------|------|
| Chlorpyrifos | 0 | | 0.00 | 0.00 | 0.00 | 0 | 0 |
| DDT | 48 | | 0.3 | 0.1 | 0.0 | 44 | 60 |
| Heptachlor | 0 | | 0.00 | 0.02 | 0.00 | 17 | 28 |
| Hexachlorobenzene | 5 | 0 | 0.03 | 0.00 | 0.00 | 6 | 7 |
| Mirex | 0.0 | | 0.000 | 0.015 | 0.000 | 0.0 | 0.0 |
| Benzo[a]pyrene | 487 | | 33 | 7 | 7 | 76 | 114 |
| Benzo[e]pyrene | 17 | | 22 | 5 | 6 | 91 | 134 |
| PCB | 1977 | 0 | 359 | 5 | 14 | 1492 | 1934 |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Menominee River



| Lege | nd | | | |
|------|----|----------|--------|-----------|
| Low | | Medium 🛑 | High 🛑 | Outlier ★ |

| Metals | Base | line | - 2009/10** | Sedime | ent - 2010 | | Tissue - 2 | 011 |
|----------------|----------|------|-------------|--------|------------|-------|------------|-----|
| (μg/g dry wt.) | Tissue 🔾 | | Sediment [| | | O, | 0 | 0 |
| Arsenic | 6.3 | • | 11.7 | 4.0 | 16.7 | 6.8 | 6.1 | |
| Cadmium | 0.35 | | 0.45 | 0.24 | 0.67 | 0.38 | 0.39 | |
| Copper | 7 | | 39 | 11 | 22 | 11 | 13 | |
| Lead | 0.9 | | 28 | 14 | 23 | 1 | 1 | |
| Methyl Mercury | 0.08 | | 0.001 | 0.001 | 0.002 | | | |
| Mercury | 0.15 | | 0.180 | 0.087 | 0.251 | 0.088 | 0.103 | |
| Zinc | 61 | 0 | 100 | 52 | 94 | 58 | 60 | |

| Organics | Baseline - 2009/10** | | Sediment - 2010 | | Tissue - 2011 | | | | |
|-------------------|----------------------|---|-----------------|--|---------------|-------|-----|-----|---|
| (ng/g dry wt.)* | Tissue 🔾 | | Sediment | | | | Ο, | 0 | 0 |
| Alpha-Chlordane | 3.0 | 0 | 1.04 | | 0.66 | 0.71 | 5.1 | 3.1 | |
| Chlorpyrifos | 0 | | 0.88 | | 0.56 | 0.53 | 0 | 0 | |
| DDT | 210 | | 5.8 | | 3.7 | 3.9 | 31 | 39 | |
| Heptachlor | 0 | 0 | 0.00 | | 0.02 | 0.05 | 0 | 0 | |
| Hexachlorobenzene | 10 | | 1.31 | | 0.29 | 0.64 | 16 | 15 | |
| Mirex | 3.4 | 0 | 0.724 | | 0.468 | 0.211 | 0.0 | 0.0 | |
| Benzo[a]pyrene | 1126 | | 608 | | 485 | 296 | 170 | 761 | |
| Benzo[e]pyrene | 740 | | 489 | | 305 | 214 | 256 | 309 | |
| PCB | 644 | 0 | 53 | | 29 | 95 | 549 | 447 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Green Bay Fox River




| Lege | nd | | | |
|------|----|----------|--------|-----------|
| Low | | Medium 🛑 | High 🛑 | Outlier ★ |

| Arsenic | 3.7 | 2.2 | | 0.0 | 2.5 |
|----------------|------|-------|---|-------|-------|
| Cadmium | 0.28 | 0.64 | | 0.19 | 1.06 |
| Copper | 13 | 34 | | 14 | 62 |
| Lead | 1.3 | 46 | | 18 | 67 |
| Methyl Mercury | 0.02 | 0.002 | | 0.001 | 0.004 |
| Mercury | 0.06 | 0.764 | * | 0.219 | 1.050 |
| Zinc | 95 | 102 | | 46 | 183 |

| Alpha-Chlordane | 3.7 | | 0.00 | | 0.00 | 1.12 |
|-------------------|-------|---|-------|---|-------|-------|
| Chlorpyrifos | 0 | | 0.00 | | 0.00 | 0.00 |
| DDT | 170 | 0 | 1.8 | | 6.5 | 2.3 |
| Heptachlor | 0 | | 0.00 | | 0.00 | 0.00 |
| Hexachlorobenzene | 4 | 0 | 0.77 | | 0.00 | 0.00 |
| Mirex | 0.0 | | 0.000 | | 0.000 | 0.000 |
| Benzo[a]pyrene | 391 | 0 | 992 | | 1090 | 1109 |
| Benzo[e]pyrene | 893 | | 641 | | 536 | 696 |
| PCB | 21439 | | 1088 | * | 276 | 1507 |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Sheboygan River



| Legend | | | |
|--------|--------|--------|-----------|
| Low 🔵 | Medium | High 🛑 | Outlier 🚖 |

| ~~~ | Zan Z | |
|-----|-------|--|
| Ex. | | |
| | | |


| Metals | Baseline - 2009/10** | | | Sediment - 2010 | | Tissue - 2011 | | |
|----------------|----------------------|---|----------|-----------------|-------|---------------|-------|-------|
| (μg/g dry wt.) | Tissue 🔾 | | Sediment | | | O, | 0 | 0 |
| Arsenic | 3.7 | | 2.0 | 1.4 | 2.1 | 4.2 | 3.7 | 4.3 |
| Cadmium | 0.62 | | 0.31 | 0.29 | 0.25 | 0.41 | 0.37 | 0.36 |
| Copper | 31 | | 25 🧧 | 27 | 31 | 15 | 9 | 11 |
| Lead | 1.8 | | 22 | 46 | 35 | 1 | 1 | 1 |
| Methyl Mercury | 0.06 | | 0.002 | 0.001 | 0.001 | | | |
| Mercury | 0.08 | • | 0.062 | 0.057 | 0.153 | 0.051 | 0.059 | 0.053 |
| Zinc | 71 | • | 73 | 78 | 93 | 57 | 46 | 54 |

| Organics | Baseline - 2009/10** | | | | Sediment - 2010 | | Tissue - 2011 | | | |
|-------------------|----------------------|---|----------|--|-----------------|-------|---------------|-------|-------|--|
| (ng/g dry wt.)* | Tissue 🔾 | | Sediment | | | | Ο, | 0 | 0 | |
| Alpha-Chlordane | 18.2 | 0 | 1.45 | | 0.00 | 0.00 | 27.5 | 33.4 | 29.1 | |
| Chlorpyrifos | 7 | | 0.00 | | 0.00 | 0.00 | 0 | 0 | 0 | |
| DDT | 438 | | 7.6 | | 21.5 | 24.4 | 680 | 745 | 919 | |
| Heptachlor | 0 | | 0.00 | | 0.00 | 0.00 | 0 | 39 | 0 | |
| Hexachlorobenzene | 15 | • | 2.17 | | 0.24 | 0.21 | 32 | 121 | 18 | |
| Mirex | 0.0 | | 0.000 | | 0.000 | 0.000 | 0.0 | 0.0 | 0.0 | |
| Benzo[a]pyrene | 419 | 0 | 99 | | 384 | 413 | 919 | 217 | 1355 | |
| Benzo[e]pyrene | 378 | • | 84 | | 300 | 287 | 534 | 392 | 911 | |
| PCB | 21969 | | 397 | | 786 | 786 | 30310 | 22273 | 38261 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

64

Milwaukee Bay



Legend

| Low | Medium 🛑 | High 🛑 | Outlier 🚖 |
|-----|----------|--------|-----------|



| Arsenic | 5.2 | | 1.7 | 3.2 | 1.7 |
|----------------|------|---|-------|-------|-------|
| Cadmium | 0.87 | 0 | 0.70 | 1.44 | 0.68 |
| Copper | 43 | • | 13 | 60 | 26 |
| Lead | 3.8 | 0 | 21 | 77 | 112 |
| Methyl Mercury | 0.02 | | 0.000 | 0.001 | 0.000 |
| Mercury | 0.03 | | 0.068 | 0.174 | 0.094 |
| Zinc | 81 | | 90 | 278 | 146 |

| Alpha-Chlordane | 46.8 | 0 | 0.47 | 3.55 | 1.62 |
|-------------------|-------|---|-------|-------|-------|
| Chlorpyrifos | 13 | • | 0.00 | 0.00 | 0.00 |
| DDT | 1018 | 0 | 9.6 | 78.0 | 44.6 |
| Heptachlor | 6 | | 0.00 | 6.08 | 0.00 |
| Hexachlorobenzene | 213 | | 7.17 | 0.557 | 0.319 |
| Mirex | 0.0 | | 0.000 | 0.000 | 0.000 |
| Benzo[a]pyrene | 1620 | | 965 | 961 | 1021 |
| Benzo[e]pyrene | 3943 | * | 597 | 743 | 607 |
| PCB | 14752 | | 152 | 661 | 602 |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Waukegan Harbor




| Lege | nd | | | |
|------|----|----------|--------|-----------|
| Low | | Medium 🛑 | High 🛑 | Outlier ★ |

| Metals | Base | line | - 2009/10** | Sediment - 2010 | ' | Tissue - 20 | 011 |
|----------------|----------|------|-------------|-----------------|-------|-------------|-------|
| (μg/g dry wt.) | Tissue 🔵 | | Sediment | | O, | 0 | 0 |
| Arsenic | 6.9 | | 10.0 | | 7.6 | 2.6 | 2.5 |
| Cadmium | 2.01 | | 3.88 | | 2.35 | 0.52 | 0.49 |
| Copper | 52 | | 85 | | 24 | 19 | 51 |
| Lead | 4.4 | | 54 | | 2 | 1 | 1 |
| Methyl Mercury | 0.01 | | 0.003 | | | | |
| Mercury | 0.04 | | 0.141 | | 0.033 | 0.036 | 0.034 |
| Zinc | 80 | • | 165 | | 68 | 51 | 57 |

| Organics | Base | line · | - 2009/10** | | Sedim | nent - 2010 | т | issue - 20 1 | 11 |
|-------------------|----------|--------|-------------|---|-------|-------------|------|---------------------|------|
| (ng/g dry wt.)* | Tissue 🔾 | | Sediment | | | | O, | 0 | 0 |
| Alpha-Chlordane | 16.2 | 0 | 0.00 | | | | 18.6 | 23.6 | 24.4 |
| Chlorpyrifos | 29 | | 0.00 | | | | 0 | 0 | 0 |
| DDT | 481 | | 0.0 | | | | 515 | 395 | 697 |
| Heptachlor | 75 | • | 0.00 | | | | 0 | 0 | |
| Hexachlorobenzene | 6 | 0 | 1.45 | | | | 9 | 13 | 9 |
| Mirex | 2.1 | 0 | 0.000 | | | | 1.7 | 1.0 | 1.2 |
| Benzo[a]pyrene | 940 | | 674 | | | | 1614 | 975 | 1209 |
| Benzo[e]pyrene | 899 | | 565 | | | | 1007 | 1143 | 1506 |
| PCB | 3924 | | 8993 | * | | | 9247 | 6154 | 5315 |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Calumet Breakwater




| Lege | nd | | | |
|------|----|--------|------|-----------|
| Low | | Medium | High | Outlier ★ |

| Arsenic | 7.3 | • | 5.5 | |
|----------------|------|---|-------|--|
| Cadmium | 3.08 | | 0.45 | |
| Copper | 28 | | 28 | |
| Lead | 2.2 | | 35 | |
| Methyl Mercury | 0.01 | | 0.001 | |
| Mercury | 0.04 | | 0.069 | |
| Zinc | 67 | | 106 | |

| Alpha-Chlordane | 9.2 | 0.57 | |
|-------------------|------|-------|--|
| Chlorpyrifos | 10 | 0.58 | |
| DDT | 144 | 19.8 | |
| Heptachlor | 0 | 0.64 | |
| Hexachlorobenzene | 4 | 0.81 | |
| Mirex | 4.0 | 0.000 | |
| Benzo[a]pyrene | 1580 | 308 | |
| Benzo[e]pyrene | 853 | 214 | |
| PCB | 1167 | 66 | |


^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Hammond Marina



| Base | line | - 2009/10** | |
|----------|---|-------------|--|
| Tissue 🔾 | | Sediment | |
| 0.0 | | 0.00 | |
| 15 | • | 0.00 | |
| 109 | 0 | 2.2 | |
| 0 | | 0.00 | |
| 5 | 0 | 0.00 | |
| 2.4 | 0 | 0.000 | |
| 779 | | 16 | |
| 805 | | 15 | |
| 1270 | 0 | 8 | |
| | 0.0 15 109 0 5 2.4 779 805 | 0.0 | 0.0 0.00 15 0.00 109 2.2 0 0.00 5 0.00 2.4 0.000 779 16 805 15 |

Kalamazoo River



| Legend | | | | | | | |
|--------|--|----------|------|--|-----------|--|--|
| Low | | Medium 🥚 | High | | Outlier ★ | | |


| ~~ | 200 | | |
|-----|-----|---------|---|
| Ex. | | A Serve | |
| - | 4 | | |
| | | | 3 |

| Arsenic | 3.7 | 5.3 |
|----------------|-------|-------|
| Cadmium | 0.28 | 0.34 |
| Copper | 10 | 11 |
| Lead | 21 | 27 |
| Methyl Mercury | 0.000 | 0.001 |
| Mercury | 0.097 | 0.191 |
| Zinc | 46 | 54 |

| Alpha-Chlordane | 0.25 | 0.42 |
|-------------------|-------|-------|
| Chlorpyrifos | 0.29 | 0.11 |
| DDT | 4.9 | 9.7 |
| Heptachlor | 0.02 | 0.09 |
| Hexachlorobenzene | 0.13 | 0.062 |
| Mirex | 0.000 | 0.000 |
| Benzo[a]pyrene | 32 | 28 |
| Benzo[e]pyrene | 23 | 24 |
| PCB | 80 | 173 |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Muskegon Lake



| Organics | Base | line | - 2009/10** | S | ec | ediment - 2010 | ediment - 2010 | ediment - 2010 Tissue - |
|-------------------|----------|------|-------------|---|----|----------------|----------------|-------------------------|
| (ng/g dry wt.)* | Tissue 🔾 | | Sediment [| | | | . O, | O, O |
| Alpha-Chlordane | 17.1 | | 1.39 | | | | | |
| Chlorpyrifos | 15 | • | 0.73 | | | | | |
| DDT | 110 | 0 | 37.3 | | | | | |
| Heptachlor | 0 | 0 | 1.19 | | | | | |
| Hexachlorobenzene | 14 | • | 0.72 | | | | | |
| Mirex | 3.8 | 0 | 0.026 | | | | | |
| Benzo[a]pyrene | 753 | | 371 | | | | | |
| Benzo[e]pyrene | 475 | 0 | 306 | | | | | |
| PCB | 773 | | 110 | | | | | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites). 70

White Lake



Low ● Medium ● Outlier ★


| ~ | approx. | |
|-----|--------------|-----|
| The | Francis, | |
| | A John Marie | - |
| - | | F 1 |
| | | 3 |

| Arsenic | 5.5 | | 3.4 | 2.1 |
|----------------|------|---|-------|-------|
| Cadmium | 0.76 | | 0.49 | 0.00 |
| Copper | 31 | | 15 | 2 |
| Lead | 1.2 | | 18 | 11 |
| Methyl Mercury | 0.05 | • | 0.003 | 0.000 |
| Mercury | 0.07 | | 0.091 | 0.015 |
| Zinc | 68 | | 64 | 36 |

| Alpha-Chlordane | 7.5 | 0.12 | 0.03 |
|-------------------|------|-------|-------|
| Chlorpyrifos | 9 | 0.27 | 0.05 |
| DDT | 89 | 6.8 | 0.2 |
| Heptachlor | 0 | 0.05 | 0.00 |
| Hexachlorobenzene | 8 | 0.12 | 0.127 |
| Mirex | 55.0 | 0.032 | 0.017 |
| Benzo[a]pyrene | 119 | 46 | 1 |
| Benzo[e]pyrene | 107 | 38 | 1 |
| PCB | 677 | 4 | 3 |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).


Saginaw River



| Organics | Baseline - 2009/10** | | | | Sediment - 2010 | | Tissue - 2011 | | |
|-------------------|----------------------|---|----------|---|-----------------|--|---------------|---|---|
| (ng/g dry wt.)* | Tissue 🔾 | | Sediment | | 3 | | O, | 0 | 0 |
| Alpha-Chlordane | 16.7 | 0 | 0.05 | | 0.18 | | | | |
| Chlorpyrifos | 11 | • | 0.05 | | 0.31 | | | | |
| DDT | 238 | 0 | 5.3 | | 11.4 | | | | |
| Heptachlor | 0 | | 0.26 | | 0.30 | | | | |
| Hexachlorobenzene | 8 | 0 | 3.86 | | 6.394 | | | | |
| Mirex | 0.0 | 0 | 0.024 | | 0.031 | | | | |
| Benzo[a]pyrene | 552 | • | 162 |] | 238 | | | | |
| Benzo[e]pyrene | 462 | • | 128 | | 205 | | | | |
| PCB | 3035 | 0 | 98 | | 90 | | | | |

^{72 *}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

St. Clair River



| Legend | | | |
|--------|----------|--------|-----------|
| Low | Medium 🛑 | High 🛑 | Outlier 🚖 |


| -way | 200 |
|------|-------------|
| 4 | Church L |
| 1 | |
| | of the said |
| 6 | |
| 5 | |

| Arsenic | 5.4 | 2.3 | 3.5 |
|----------------|------|-------|-------|
| Cadmium | 3.55 | 0.00 | 0.49 |
| Copper | 80 | 5 | 21 |
| Lead | 3.1 | 9 | 15 |
| Methyl Mercury | 0.02 | 0.001 | 0.002 |
| Mercury | 0.06 | 0.012 | 0.138 |
| Zinc | 135 | 28 | 62 |

| Alaba Oblandana | 0.0 | 0.00 | _ | 0.00 |
|-------------------|-----|-------|---|--------|
| Alpha-Chlordane | 8.0 | 0.00 | | 0.02 |
| Chlorpyrifos | 0 | 0.06 | | 0.12 |
| DDT | 24 | 0.3 | | 1.7 |
| Heptachlor | 0 | 0.00 | | 0.10 |
| Hexachlorobenzene | 2 | 0.57 | | 10.857 |
| Mirex | 1.2 | 0.000 | | 0.094 |
| Benzo[a]pyrene | 0 | 6 | | 33 |
| Benzo[e]pyrene | 0 | 5 | | 25 |
| PCB | 173 | 2 | | 24 |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Clinton River



| Organics | Base | line | - 2009/10** | | Sedimer | nt - 2010 | | Tissue - | 2011 |
|-------------------|----------|------|-------------|---|---------|-----------|----|----------|------|
| (ng/g dry wt.)* | Tissue 🔾 | | Sediment | | | | O, | 0 | 0 |
| Alpha-Chlordane | 43.8 | | 4.66 | | 0.46 | | | | |
| Chlorpyrifos | 22 | | 0.61 | | 0.55 | | | | |
| DDT | 474 | | 68.1 | * | 66.9 | | | | |
| Heptachlor | 0 | | 1.07 | | 1.54 | | | | |
| Hexachlorobenzene | 7 | | 1.77 | | 1.257 | | | | |
| Mirex | 1.5 | | 0.552 | | 0.422 | | | | |
| Benzo[a]pyrene | 169 | 0 | 3569 | * | 2594 | | | | |
| Benzo[e]pyrene | 1038 | | 3369 | * | 1988 | | | | |
| PCB | 2636 | 0 | 367 | | 378 | | | | |

0.204

380

0.07

214

0.182

330

Mercury

Zinc

^{74 *}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Rouge River



| Lege | nd | | | |
|------|----|----------------|--------|-----------|
| Low | | Medium <u></u> | High 🛑 | Outlier 🚖 |


| ~~ | -05 | | |
|-----|--|--|---|
| Ta | Runi | Z. | |
| { } | The same of the sa | The state of the s | |
| | _{} | | 5 |
| | | | |

| Arsenic | 4.0 | |
|----------------|-------|--|
| Cadmium | 0.86 | |
| Copper | 94 | |
| Lead | 54 | |
| Methyl Mercury | 0.001 | |
| Mercury | 0.195 | |
| Zinc | 180 | |

| Alpha-Chlordane | 13.10 | * |
|-------------------|-------|---|
| Chlorpyrifos | 0.00 | |
| DDT | 58.6 | * |
| Heptachlor | 9.96 | |
| Hexachlorobenzene | 0.27 | |
| Mirex | 0.062 | |
| Benzo[a]pyrene | 16454 | * |
| Benzo[e]pyrene | 6715 | * |
| PCB | 372 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Detroit River



| Organics | Base | line | - 2009/10** | Sediment - 2010 | 1 | Tissue - 20 | 011 |
|-------------------|----------|------|-------------|-----------------|------|-------------|-----|
| (ng/g dry wt.)* | Tissue 🔾 | | Sediment | | O, | 0 | 0 |
| Alpha-Chlordane | 7.4 | 0 | 0.67 | | 0.0 |) | |
| Chlorpyrifos | 9 | • | 0.00 | | 0 |) | |
| DDT | 73 | | 9.2 | | 454 | | |
| Heptachlor | 0 | | 0.00 | | 0 |) | |
| Hexachlorobenzene | 12 | 0 | 4.08 | | 16 | ; | |
| Mirex | 0.0 | | 0.000 | | 0.0 |) | |
| Benzo[a]pyrene | 246 | 0 | 401 | | 1852 | ! | |
| Benzo[e]pyrene | 441 | • | 288 | | 2602 | | |
| PCB | 1652 | | 68 | | 5126 | i | |

114

86

46

Zinc

76

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

River Raisin



| Leger | nd | | | |
|-------|----|--------|------|-----------|
| Low | | Medium | High | Outlier ★ |


| ~~ | - | | |
|-----|--|---|---|
| The | Runi | 2 | |
| [[| The state of the s | | |
| -}_ | 1 | E | 5 |
| | | | |

| Arsenic | 3.8 | 6.6 | |
|----------------|------|-------|--|
| Cadmium | 1.97 | 0.60 | |
| Copper | 16 | 26 | |
| Lead | 2.6 | 29 | |
| Methyl Mercury | 0.01 | 0.002 | |
| Mercury | 0.04 | 0.077 | |
| Zinc | 54 | 94 | |

| Alpha-Chlordane | 33.5 | • | 0.97 | |
|-------------------|------|---|-------|--|
| Chlorpyrifos | 32 | | 0.00 | |
| DDT | 225 | | 21.2 | |
| Heptachlor | 0 | | 0.00 | |
| Hexachlorobenzene | 16 | | 0.38 | |
| Mirex | 1.5 | | 0.000 | |
| Benzo[a]pyrene | 288 | | 1210 | |
| Benzo[e]pyrene | 578 | • | 616 | |
| PCB | 4562 | • | 223 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Maumee River




| Organics | Base | line | - 2009/10** | |
|-------------------|----------|------|-------------|--|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment [| |
| Alpha-Chlordane | 48.2 | | 0.69 | |
| Chlorpyrifos | 249 | * | 0.53 | |
| DDT | 216 | 0 | 7.9 | |
| Heptachlor | 0 | | 0.21 | |
| Hexachlorobenzene | 7 | | 0.19 | |
| Mirex | 1.6 | | 0.000 | |
| Benzo[a]pyrene | 543 | 0 | 190 | |
| Benzo[e]pyrene | 1095 | | 153 | |
| PCB | 2726 | | 101 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).


78

Black River



^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Cuyahoga River



| Metals | Base | Baseline - 2009/10** | | Sediment - 2010 | | Tissue - 2011 | | |
|----------------|----------|----------------------|----------|-----------------|--|---------------|-------|-------|
| (μg/g dry wt.) | Tissue 🔾 | | Sediment | | | O, | 0 | 0 |
| Arsenic | 4.3 | 0 | 17.1 | | | 5.7 | 5.8 | 5.2 |
| Cadmium | 0.95 | | 0.93 | | | 1.60 | 1.55 | 1.31 |
| Copper | 50 | • | 43 | | | 20 | 54 | 38 |
| Lead | 4.4 | | 41 | | | 3 | 5 | 4 |
| Methyl Mercury | 0.02 | | 0.001 | | | | | |
| Mercury | 0.03 | | 0.095 | | | 0.044 | 0.048 | 0.050 |
| Zinc | 76 | • | 203 | | | 69 | 90 | 79 |

| Organics | Base | Baseline - 2009/10** | | Sediment - 2010 | Tissue - 201 | | 11 | |
|-------------------|----------|----------------------|----------|-----------------|--------------|-------|-------|-------|
| (ng/g dry wt.)* | Tissue 🔾 | | Sediment | | | O, | 0 | 0 |
| Alpha-Chlordane | 177.3 | | 3.14 | | | 195.9 | 225.0 | 218.3 |
| Chlorpyrifos | 23 | | 0.00 | | | 0 | 0 | 0 |
| DDT | 344 | | 13.9 | | | 516 | 596 | 602 |
| Heptachlor | 0 | | 0.00 | | | 0 | 0 | 0 |
| Hexachlorobenzene | 10 | 0 | 1.51 | | | 0 | 0 | 0 |
| Mirex | 3.5 | | 0.000 | | | 0.0 | 0.0 | 0.0 |
| Benzo[a]pyrene | 756 | 0 | 454 | | | 1080 | 1245 | 540 |
| Benzo[e]pyrene | 2087 | | 418 | | | 2980 | 3286 | 1660 |
| PCB | 3557 | • | 163 | | | 3903 | 4664 | 4161 |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Ashtabula River



| Legend | | | |
|--------|----------|--------|-----------|
| Low | Medium 🛑 | High 🛑 | Outlier ★ |


| 1 | |
|---|--|

| Metals | Base | Baseline - 2009/10** | | | Sedim | ent - 2010 | Tissue - 2011 | | |
|-------------------------|----------|----------------------|----------|--|-------|------------|---------------|-------|-------|
| (μ g/g dry wt.) | Tissue 🔾 | | Sediment | | | | Ο, | 0 | 0 |
| Arsenic | 7.9 | | 14.2 | | | | 5.1 | 6.6 | 6.3 |
| Cadmium | 2.34 | | 0.59 | | | | 2.74 | 2.71 | 1.77 |
| Copper | 124 | | 30 | | | | 14 | 15 | 15 |
| Lead | 5.4 | | 25 | | | | 1 | 1 | 2 |
| Methyl Mercury | 0.02 | | 0.001 | | | | | | |
| Mercury | 0.04 | | 0.055 | | | | 0.038 | 0.045 | 0.045 |
| Zinc | 97 | | 117 | | | | 50 | 60 | 59 |

| Organics | Base | Baseline - 2009/10** | | Sedim | nent - 2010 | т | issue - 20 | 11 | |
|-------------------|----------|----------------------|------------|-------|-------------|---|------------|------|------|
| (ng/g dry wt.)* | Tissue 🔾 | | Sediment [| | | | O, | 0 | 0 |
| Alpha-Chlordane | 19.2 | 0 | 3.83 | | | | 9.2 | 23.4 | 10.7 |
| Chlorpyrifos | 6 | 0 | 0.00 | | | | 0 | 0 | 0 |
| DDT | 46 | | 7.1 | | | | 62 | 99 | 50 |
| Heptachlor | 0 | | 0.00 | | | | 0 | 0 | 0 |
| Hexachlorobenzene | 131 | | 4.99 | | | | 58 | 137 | 130 |
| Mirex | 0.0 | | 0.000 | | | | 0.0 | 8.3 | 0.0 |
| Benzo[a]pyrene | 403 | 0 | 361 | | | | 94 | 145 | 323 |
| Benzo[e]pyrene | 627 | • | 250 | | | | 208 | 360 | 640 |
| PCB | 1289 | 0 | 102 | | | | 1067 | 1979 | 1292 |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Presque Isle Bay



| Organics | Baseline - 2009/10** | | | |
|-------------------|----------------------|---|----------|---|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment | |
| Alpha-Chlordane | 84.7 | | 2.96 | |
| Chlorpyrifos | 4 | | 0.00 | |
| DDT | 224 | 0 | 7.2 | |
| Heptachlor | 0 | 0 | 0.00 | |
| Hexachlorobenzene | 238 | | 8.56 | |
| Mirex | 22.6 | • | 0.000 | |
| Benzo[a]pyrene | 38041 | * | 2380 | * |
| Benzo[e]pyrene | 23196 | * | 1300 | * |
| PCB | 4138 | | 182 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Buffalo River



Legend

Low Medium High Outlier 🛨




| Arsenic | 4.4 | • | 13.1 | |
|----------------|------|---|-------|--|
| Cadmium | 0.78 | • | 0.50 | |
| Copper | 43 | • | 41 | |
| Lead | 5.3 | | 32 | |
| Methyl Mercury | 0.08 | | 0.002 | |
| Mercury | 0.11 | | 0.076 | |
| Zinc | 82 | • | 139 | |

| Alpha-Chlordane | 24.4 | | 1.22 | |
|-------------------|------|---|-------|--|
| Chlorpyrifos | 10 | • | 0.39 | |
| DDT | 249 | | 7.8 | |
| Heptachlor | 0 | | 0.00 | |
| Hexachlorobenzene | 17 | • | 0.69 | |
| Mirex | 4.5 | | 0.000 | |
| Benzo[a]pyrene | 879 | • | 201 | |
| Benzo[e]pyrene | 1313 | | 179 | |
| PCB | 1789 | • | 53 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Niagara River



| Organics | Base | Baseline - 2009/10** | | Sediment - 2010 | liment - 2010 Tissue - 201 | | 2011 | |
|-------------------|----------|----------------------|----------|-----------------|----------------------------|----|------|---|
| (ng/g dry wt.)* | Tissue 🔾 | | Sediment | l | | O, | 0 | 0 |
| Alpha-Chlordane | 12.9 | | 0.16 | | | 9 | 9.3 | |
| Chlorpyrifos | 5 | | 0.00 | | | | 0 | |
| DDT | 84 | 0 | 1.0 | | | 1 | 01 | |
| Heptachlor | 0 | 0 | 0.00 | | | | 0 | |
| Hexachlorobenzene | 25 | | 0.58 | | | | 0 | |
| Mirex | 6.3 | 0 | 0.000 | | | (| 0.0 | |
| Benzo[a]pyrene | 1067 | | 26 | | | 24 | 43 | |
| Benzo[e]pyrene | 1416 | | 28 | | | 40 | 29 | |
| PCB | 4715 | | 25 | | | 50 | 64 | |

57


89

51

Zinc

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Eighteenmile Creek



| Lege | nd | | | |
|------|----|----------|--------|-----------|
| Low | | Medium 🛑 | High 🛑 | Outlier ★ |


| The state of the s | | |
|--|--|--|
| | | |

| Arsenic | 2.9 | • | 4.0 | |
|----------------|------|---|-------|--|
| Cadmium | 0.70 | | 0.48 | |
| Copper | 56 | • | 63 | |
| Lead | 9.3 | | 47 | |
| Methyl Mercury | 0.05 | • | 0.002 | |
| Mercury | 0.08 | | 0.127 | |
| Zinc | 96 | | 182 | |

| Alpha-Chlordane | 37.7 | • | 0.77 | |
|-------------------|-------|---|-------|--|
| Chlorpyrifos | 15 | • | 0.36 | |
| DDT | 1896 | | 21.1 | |
| Heptachlor | 0 | | 0.00 | |
| Hexachlorobenzene | 22 | • | 0.60 | |
| Mirex | 52.0 | | 0.310 | |
| Benzo[a]pyrene | 344 | • | 170 | |
| Benzo[e]pyrene | 791 | | 135 | |
| PCB | 23964 | | 170 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Rochester



| Organics | Base | line | - 2009/10** | | Sediment - 2010 | | Tissue - | 2011 |
|-------------------|----------|------|-------------|---|-----------------|----|----------|------|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment | l | - | O, | 0 | 0 |
| Alpha-Chlordane | 10.9 | 0 | 0.64 | | | 10 | 0.0 | |
| Chlorpyrifos | 8 | • | 0.30 | | | | 0 | |
| DDT | 119 | | 6.9 | | | 2 | 16 | |
| Heptachlor | 0 | | 0.06 | | | | 0 | |
| Hexachlorobenzene | 16 | | 3.23 | | | | 0 | |
| Mirex | 42.8 | | 0.000 | | | C | 0.0 | |
| Benzo[a]pyrene | 173 | | 147 | | | 19 | 97 | |
| Benzo[e]pyrene | 395 | • | 113 | | | 3 | 32 | |
| PCB | 681 | | 32 | | | 4 | 34 | |

Oswego River




Legend Low ● Medium ● High ● Outlier ★

| Arsenic | 3.6 | | 1.7 | |
|----------------|------|---|-------|--|
| Cadmium | 0.65 | | 0.16 | |
| Copper | 23 | | 9 | |
| Lead | 0.9 | | 15 | |
| Methyl Mercury | 0.04 | • | 0.001 | |
| Mercury | 0.06 | 0 | 0.065 | |
| Zinc | 57 | | 36 | |

| Alpha-Chlordane | 5.2 | | 0.33 | |
|-------------------|------|---|-------|--|
| Chlorpyrifos | 4 | 0 | 0.08 | |
| DDT | 64 | 0 | 2.9 | |
| Heptachlor | 0 | 0 | 0.00 | |
| Hexachlorobenzene | 21 | | 0.38 | |
| Mirex | 16.3 | • | 1.310 | |
| Benzo[a]pyrene | 106 | 0 | 213 | |
| Benzo[e]pyrene | 380 | • | 140 | |
| PCB | 667 | | 17 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).


St. Lawrence River




| Base | line | - 2009/10** | |
|----------|---|-------------|-----|
| Tissue 🔾 | | Sediment | |
| 6.0 | | 0.04 | |
| 14 | • | 0.00 | |
| 40 | 0 | 0.5 | |
| 0 | 0 | 0.02 | |
| 4 | 0 | 0.17 | |
| 22.7 | • | 0.000 | |
| 386 | | 12 | |
| 621 | • | 12 | |
| 399 | 0 | 5 | |
| | 6.0 14 40 0 4 22.7 386 621 | 6.0 | 6.0 |




Reference sites





Bayshore Park




Low Medium High Outlier ★

| Metals | Baseline** | | | | |
|----------------|------------|---|----------|--|--|
| (μg/g dry wt.) | Tissue 🔾 | | Sediment | | |
| Arsenic | 3.7 | 0 | 1.6 | | |
| Cadmium | 0.35 | | 0.39 | | |
| Copper | 206 | | 13 | | |
| Lead | 7.0 | | 21 | | |
| Methyl Mercury | 0.06 | | 0.001 | | |
| Mercury | 0.08 | | 0.261 | | |
| Zinc | 200 | * | 53 | | |

| Organics | Baseline** | | | |
|-------------------|------------|---|------------|--|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment [| |
| Alpha-Chlordane | 3.3 | 0 | 0.28 | |
| Chlorpyrifos | 2 | | 0.65 | |
| DDT | 109 | | 1.3 | |
| Heptachlor | 53 | • | 0.04 | |
| Hexachlorobenzene | 18 | 0 | 0.26 | |
| Mirex | 0.9 | | 0.000 | |
| Benzo[a]pyrene | 241 | 0 | 189 | |
| Benzo[e]pyrene | 151 | | 115 | |
| PCB | 4741 | | 100 | |

North Chicago



Legend

Low Medium High Outlier 🛨



| Metals | Baseline** | | | | |
|------------------------|------------|----------|--|--|--|
| (μ g/g dry wt.) | Tissue 🔵 | Sediment | | | |
| Arsenic | 6.2 | • | | | |
| Cadmium | 3.01 | • | | | |
| Copper | 28 | | | | |
| Lead | 8.5 | | | | |
| Methyl Mercury | 0.01 | | | | |
| Mercury | 0.03 | | | | |
| Zinc | 74 | • | | | |

| Organics | Baseline** | | | |
|-------------------|------------|----------|--|--|
| (ng/g dry wt.)* | Tissue 🔵 | Sediment | | |
| Alpha-Chlordane | 26.0 | • | | |
| Chlorpyrifos | 22 | • | | |
| DDT | 788 | | | |
| Heptachlor | 0 | | | |
| Hexachlorobenzene | 6 | | | |
| Mirex | 4.1 | | | |
| Benzo[a]pyrene | 794 | • | | |
| Benzo[e]pyrene | 562 | • | | |
| PCB | 1851 | | | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Holland Breakwater



Legend Low ● Medium ● High ● Outlier ★



| Metals | | Baseline** |
|------------------------|----------|------------|
| (μ g/g dry wt.) | Tissue 🔵 | Sediment |
| Arsenic | 5.0 | • |
| Cadmium | 1.92 | • |
| Copper | 28 | |
| Lead | 1.2 | |
| Methyl Mercury | 0.03 | |
| Mercury | 0.05 | |
| Zinc | 75 | |

| Organics | 1 | Base | eline** |
|-------------------|----------|------|-------------------|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment ■ |
| Alpha-Chlordane | 19.3 | 0 | |
| Chlorpyrifos | 10 | | |
| DDT | 211 | | |
| Heptachlor | 0 | | |
| Hexachlorobenzene | 9 | | |
| Mirex | 0.0 | | |
| Benzo[a]pyrene | 187 | | |
| Benzo[e]pyrene | 142 | | |
| PCB | 1192 | | |

Muskegon



Legend Low Medium Outlier *

| Metals | | Baseline** |
|----------------|----------|------------|
| (μg/g dry wt.) | Tissue 🔵 | Sediment |
| Arsenic | 4.3 | |
| Cadmium | 1.39 | |
| Copper | 13 | |
| Lead | 1.2 | |
| Methyl Mercury | 0.02 | |
| Mercury | 0.04 | |
| Zinc | 53 | |

| Organics | Baseline** | | |
|-------------------|------------|----------|--|
| (ng/g dry wt.)* | Tissue 🔵 | Sediment | |
| Alpha-Chlordane | 18.9 | • | |
| Chlorpyrifos | 14 | • | |
| DDT | 145 | | |
| Heptachlor | 0 | | |
| Hexachlorobenzene | 6 | | |
| Mirex | 3.5 | | |
| Benzo[a]pyrene | 275 | | |
| Benzo[e]pyrene | 114 | | |
| PCB | 665 | | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Leelanau State Park




Legend Low ● Medium ● High ● Outlier ★



| Metals | | Baseline** |
|----------------|----------|-------------------|
| (μg/g dry wt.) | Tissue 🔵 | Sediment = |
| Arsenic | 9.9 | • |
| Cadmium | 4.24 | • |
| Copper | 44 | • |
| Lead | 1.6 | |
| Methyl Mercury | 0.02 | |
| Mercury | 0.05 | |
| Zinc | 69 | • |

| Organics | Baseline** | | |
|-------------------|------------|----------|--|
| (ng/g dry wt.)* | Tissue 🔵 | Sediment | |
| Alpha-Chlordane | 5.3 | • | |
| Chlorpyrifos | 6 | | |
| DDT | 15 | | |
| Heptachlor | 0 | | |
| Hexachlorobenzene | 0 | | |
| Mirex | 0.6 | | |
| Benzo[a]pyrene | 0 | | |
| Benzo[e]pyrene | 33 | | |
| PCB | 103 | | |

Lincoln Bay




Low ● Medium ● Outlier ★

| Metals | | Baseline** |
|----------------|----------|------------|
| (μg/g dry wt.) | Tissue 🔵 | Sediment |
| Arsenic | 9.5 | • |
| Cadmium | 6.93 | • |
| Copper | 19 | |
| Lead | 0.4 | |
| Methyl Mercury | 0.03 | |
| Mercury | 0.09 | • |
| Zinc | 68 | |

| Organics | Baseline** | | |
|-------------------|------------|----------|--|
| (ng/g dry wt.)* | Tissue 🔵 | Sediment | |
| Alpha-Chlordane | 3.4 | • | |
| Chlorpyrifos | 0 | | |
| DDT | 14 | | |
| Heptachlor | 0 | | |
| Hexachlorobenzene | 3 | | |
| Mirex | 0.9 | | |
| Benzo[a]pyrene | 0 | | |
| Benzo[e]pyrene | 0 | | |
| PCB | 75 | | |


Thunder Bay



| Metals | Baseline** | | | |
|------------------------|------------|---|----------|--|
| (μ g/g dry wt.) | Tissue 🔵 | | Sediment | |
| Arsenic | 5.4 | | 1.3 | |
| Cadmium | 8.44 | | 0.24 | |
| Copper | 125 | | 9 | |
| Lead | 4.9 | 0 | 10 | |
| Methyl Mercury | 0.02 | | 0.002 | |
| Mercury | 0.07 | 0 | 0.040 | |
| Zinc | 139 | | 34 | |

| Organics | Baseline** | | | |
|-------------------|------------|---|------------|--|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment [| |
| Alpha-Chlordane | 9.5 | 0 | 0.07 | |
| Chlorpyrifos | 3 | | 0.00 | |
| DDT | 30 | | 0.6 | |
| Heptachlor | 0 | | 0.00 | |
| Hexachlorobenzene | 5 | | 0.098 | |
| Mirex | 0.0 | | 0.011 | |
| Benzo[a]pyrene | 0 | | 85 | |
| Benzo[e]pyrene | 0 | | 46 | |
| PCB | 75 | | 2 | |

Eagle Bay



Legend Low ● Medium ● High ● Outlier ★

| | San | |
|--|---|--|
| | | |

| Metals | Baseline** | |
|----------------|------------|----------|
| (μg/g dry wt.) | Tissue 🔵 | Sediment |
| Arsenic | 6.7 | • |
| Cadmium | 3.94 | |
| Copper | 51 | |
| Lead | 0.9 | |
| Methyl Mercury | 0.02 | |
| Mercury | 0.06 | |
| Zinc | 109 | |

| Organics | Baseline** | | |
|-------------------|------------|---|----------|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment |
| Alpha-Chlordane | 8.3 | 0 | |
| Chlorpyrifos | 12 | 0 | |
| DDT | 38 | | |
| Heptachlor | 8 | | |
| Hexachlorobenzene | 4 | 0 | |
| Mirex | 0.0 | | |
| Benzo[a]pyrene | 625 | 0 | |
| Benzo[e]pyrene | 54 | | |
| PCB | 245 | | |

Black River Canal



Legend Low 🔵 Medium ___ High (Outlier 👚



| Metals | Baseline** | | |
|----------------|------------|-------------------|--|
| (μg/g dry wt.) | Tissue 🔵 | Sediment ■ | |
| Arsenic | 8.1 | • | |
| Cadmium | 5.57 | • | |
| Copper | 38 | • | |
| Lead | 1.5 | | |
| Methyl Mercury | 0.02 | | |
| Mercury | 0.08 | | |
| Zinc | 102 | | |

| Organics | Baseline** | | |
|-------------------|------------|----------|--|
| (ng/g dry wt.)* | Tissue 🔵 | Sediment | |
| Alpha-Chlordane | 8.7 | 0 | |
| Chlorpyrifos | 22 | | |
| DDT | 48 | 0 | |
| Heptachlor | 0 | | |
| Hexachlorobenzene | 2 | 0 | |
| Mirex | 1.3 | | |
| Benzo[a]pyrene | 0 | 0 | |
| Benzo[e]pyrene | 0 | | |
| PCB | 161 | | |

Anchor Bay



Low ● Medium ● Outlier ★

| _ | | |
|---|--|--|

| Metals | Baseline** | | |
|----------------|------------|------------|--|
| (μg/g dry wt.) | Tissue 🔵 | Sediment (| |
| Arsenic | | 3.3 | |
| Cadmium | | 0.68 | |
| Copper | | 17 | |
| Lead | | 12 | |
| Methyl Mercury | | 0.001 | |
| Mercury | | 0.072 | |
| Zinc | | 62 | |

| Organics | | Baseline** | |
|-------------------|----------|------------|--|
| (ng/g dry wt.)* | Tissue 🔵 | Sediment | |
| Alpha-Chlordane | | 0.02 | |
| Chlorpyrifos | | 0.00 | |
| DDT | | 0.8 | |
| Heptachlor | | 0.04 | |
| Hexachlorobenzene | | 4.069 | |
| Mirex | | 0.000 | |
| Benzo[a]pyrene | | 25 | |
| Benzo[e]pyrene | | 18 | |
| PCB | | 10 | |

Stony Point



| Legend | | | |
|--------|----------|--------|-----------|
| Low 🔵 | Medium 🛑 | High 🛑 | Outlier ★ |

| Metals | Baseline** | | | |
|----------------|------------|---|------------|--|
| (μg/g dry wt.) | Tissue 🔵 | | Sediment [| |
| Arsenic | 3.9 | | 4.1 | |
| Cadmium | 2.11 | 0 | 0.98 | |
| Copper | 69 | | 23 | |
| Lead | 5.3 | 0 | 26 | |
| Methyl Mercury | 0.04 | 0 | 0.001 | |
| Mercury | 0.09 | | 0.306 | |
| Zinc | 90 | | 100 | |

| Organics | Baseline** | | | |
|-------------------|------------|---|------------|--|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment [| |
| Alpha-Chlordane | 37.1 | • | 0.54 | |
| Chlorpyrifos | 18 | | 0.00 | |
| DDT | 408 | | 9.8 | |
| Heptachlor | 0 | | 1.49 | |
| Hexachlorobenzene | 20 | 0 | 1.08 | |
| Mirex | 3.5 | | 0.000 | |
| Benzo[a]pyrene | 577 | • | 174 | |
| Benzo[e]pyrene | 1293 | | 139 | |
| PCB | 7334 | | 95 | |

Reno Beach




| A Part of the second of the se | |
|--|--|
| | |

| Metals | Baseline** | | | |
|----------------|------------|------------|-------|--|
| (μg/g dry wt.) | Tissue 🔵 | Sediment ■ | | |
| Arsenic | 5.0 | | 9.0 | |
| Cadmium | 1.79 | 0 | 0.55 | |
| Copper | 109 | | 16 | |
| Lead | 4.0 | 0 | 15 | |
| Methyl Mercury | 0.02 | | 0.001 | |
| Mercury | 0.03 | | 0.057 | |
| Zinc | 92 | | 53 | |

| Organics | | Base | eline** | |
|-------------------|----------|------|----------|--|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment | |
| Alpha-Chlordane | 10.1 | 0 | 0.50 | |
| Chlorpyrifos | 17 | | 0.42 | |
| DDT | 43 | | 7.6 | |
| Heptachlor | 0 | | 0.16 | |
| Hexachlorobenzene | 15 | 0 | 0.64 | |
| Mirex | 0.0 | | 0.000 | |
| Benzo[a]pyrene | 156 | 0 | 35 | |
| Benzo[e]pyrene | 335 | • | 47 | |
| PCB | 749 | 0 | 51 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Peach Orchard Pt.



| Legend | | | |
|--------|----------|--------|-----------|
| Low 🔵 | Medium 🛑 | High 🛑 | Outlier ★ |

| A Comment of the comm | |
|--|--|
| 1 | |

| Metals | Baseline** | | | |
|----------------|------------|----------|-------|--|
| (μg/g dry wt.) | Tissue 🔵 | Sediment | | |
| Arsenic | 5.1 | 0 | 6.5 | |
| Cadmium | 2.61 | 0 | 0.96 | |
| Copper | 12 | | 30 | |
| Lead | 2.7 | | 35 | |
| Methyl Mercury | 0.01 | | 0.001 | |
| Mercury | 0.04 | | 0.179 | |
| Zinc | 54 | | 133 | |

| Organics | Baseline** | | | |
|-------------------|------------|---|------------|--|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment [| |
| Alpha-Chlordane | 15.0 | 0 | 0.73 | |
| Chlorpyrifos | 20 | | 0.24 | |
| DDT | 109 | 0 | 8.9 | |
| Heptachlor | 0 | | 0.53 | |
| Hexachlorobenzene | 9 | 0 | 0.84 | |
| Mirex | 1.9 | | 0.000 | |
| Benzo[a]pyrene | 130 | 0 | 82 | |
| Benzo[e]pyrene | 391 | • | 95 | |
| PCB | 2857 | | 90 | |

Old Woman Creek



Legend Low ● Medium ● Outlier ★



| Metals | Baseline** | | | |
|----------------|------------|----------|-------|--|
| (μg/g dry wt.) | Tissue 🔵 | Sediment | | |
| Arsenic | 4.2 | 0 | 11.8 | |
| Cadmium | 2.11 | | 0.86 | |
| Copper | 12 | | 31 | |
| Lead | 1.2 | | 26 | |
| Methyl Mercury | 0.01 | | 0.001 | |
| Mercury | 0.03 | | 0.075 | |
| Zinc | 56 | | 91 | |

| Organics | Baseline** | | | |
|-------------------|------------|---|----------|--|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment | |
| Alpha-Chlordane | 14.9 | 0 | 0.79 | |
| Chlorpyrifos | 16 | • | 0.82 | |
| DDT | 50 | | 23.9 | |
| Heptachlor | 0 | | 0.06 | |
| Hexachlorobenzene | 5 | 0 | 0.34 | |
| Mirex | 0.0 | | 0.020 | |
| Benzo[a]pyrene | 40 | 0 | 234 | |
| Benzo[e]pyrene | 77 | | 164 | |
| PCB | 373 | 0 | 30 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Lorain




Legend Low Outlier 🚖 Medium _ High (



| Metals | | Baseline** | | |
|------------------------|----------|------------|--|--|
| (μ g/g dry wt.) | Tissue 🔵 | Sediment | | |
| Arsenic | 3.9 | • | | |
| Cadmium | 2.66 | • | | |
| Copper | 10 | | | |
| Lead | 1.6 | | | |
| Methyl Mercury | 0.02 | | | |
| Mercury | 0.03 | | | |
| Zinc | 47 | | | |

| Organics | Baseline** | | |
|-------------------|------------|---|----------|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment |
| Alpha-Chlordane | 23.2 | 0 | |
| Chlorpyrifos | 11 | | |
| DDT | 58 | | |
| Heptachlor | 0 | | |
| Hexachlorobenzene | 6 | | |
| Mirex | 0.9 | | |
| Benzo[a]pyrene | 96 | | |
| Benzo[e]pyrene | 320 | • | |
| PCB | 604 | | |

Ashtabula



| Metals | Baseline** | | | |
|----------------|------------|------------|-------|--|
| (μg/g dry wt.) | Tissue 🔵 | Sediment ■ | | |
| Arsenic | 6.1 | 0 | 10.6 | |
| Cadmium | 3.52 | 0 | 0.54 | |
| Copper | 10 | | 22 | |
| Lead | 1.4 | | 23 | |
| Methyl Mercury | 0.01 | | 0.000 | |
| Mercury | 0.03 | | 0.058 | |
| Zinc | 58 | | 106 | |

| Organics | Baseline** | | | |
|-------------------|------------|---|----------|--|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment | |
| Alpha-Chlordane | 15.1 | 0 | 0.72 | |
| Chlorpyrifos | 4 | | 0.05 | |
| DDT | 49 | 0 | 9.0 | |
| Heptachlor | 0 | | 0.18 | |
| Hexachlorobenzene | 21 | 0 | 1.46 | |
| Mirex | 2.6 | | 0.000 | |
| Benzo[a]pyrene | 63 | 0 | 128 | |
| Benzo[e]pyrene | 135 | | 118 | |
| PCB | 1120 | 0 | 109 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Dunkirk




Legend Low 🔵 Medium _ High (Outlier 👚



| Metals | Baseline** | | | |
|----------------|------------|----------|-------|--|
| (μg/g dry wt.) | Tissue 🔵 | Sediment | | |
| Arsenic | 6.6 | 0 | 12.3 | |
| Cadmium | 3.99 | 0 | 0.36 | |
| Copper | 11 | | 15 | |
| Lead | 2.0 | | 22 | |
| Methyl Mercury | 0.01 | | 0.000 | |
| Mercury | 0.03 | | 0.033 | |
| Zinc | 56 | | 109 | |

| Organics | Baseline** | | | |
|-------------------|------------|---|------------|--|
| (ng/g dry wt.)* | Tissue 🔵 | | Sediment [| |
| Alpha-Chlordane | 7.5 | 0 | 0.53 | |
| Chlorpyrifos | 4 | | 0.07 | |
| DDT | 53 | 0 | 8.2 | |
| Heptachlor | 0 | | 0.00 | |
| Hexachlorobenzene | 10 | | 1.26 | |
| Mirex | 0.0 | | 0.000 | |
| Benzo[a]pyrene | 139 | 0 | 57 | |
| Benzo[e]pyrene | 219 | | 65 | |
| PCB | 599 | | 31 | |

Olcott



| A Company of the Comp | |
|--|--|
| | |

| Metals | Baseline** | | | | |
|----------------|------------|----------|-------|--|--|
| (μg/g dry wt.) | Tissue 🔵 | Sediment | | | |
| Arsenic | 6.2 | 0 | 10.6 | | |
| Cadmium | 3.59 | 0 | 0.54 | | |
| Copper | 9 | | 28 | | |
| Lead | 0.9 | | 25 | | |
| Methyl Mercury | 0.02 | | 0.002 | | |
| Mercury | 0.04 | | 0.176 | | |
| Zinc | 57 | | 122 | | |

| Organics | Baseline** | | | |
|-------------------|------------|----------|-------|--|
| (ng/g dry wt.)* | Tissue 🔵 | Sediment | | |
| Alpha-Chlordane | 9.9 | 0 | 0.81 | |
| Chlorpyrifos | 8 | • | 0.10 | |
| DDT | 117 | 0 | 7.7 | |
| Heptachlor | 0 | | 0.00 | |
| Hexachlorobenzene | 17 | 0 | 6.1 | |
| Mirex | 47.4 | | 0.400 | |
| Benzo[a]pyrene | 77 | 0 | 149 | |
| Benzo[e]pyrene | 207 | | 129 | |
| PCB | 866 | 0 | 99 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).

Oswego





| Metals | Baseline** | | |
|------------------------|------------|--|----------|
| (μ g/g dry wt.) | Tissue 🔵 | | Sediment |
| Arsenic | 3.5 | | |
| Cadmium | 1.08 | | |
| Copper | 12 | | |
| Lead | 0.5 | | |
| Methyl Mercury | 0.01 | | |
| Mercury | 0.03 | | |
| Zinc | 34 | | |

| Organics | Baseline** | | |
|-------------------|------------|-----|--------|
| (ng/g dry wt.)* | Tissue 🔵 | Sec | diment |
| Alpha-Chlordane | 9.7 | • | |
| Chlorpyrifos | 5 | | |
| DDT | 62 | | |
| Heptachlor | 0 | | |
| Hexachlorobenzene | 9 | | |
| Mirex | 18.7 | • | |
| Benzo[a]pyrene | 100 | | |
| Benzo[e]pyrene | 61 | | |
| PCB | 281 | | |

Cape Vincent



| Metals | Baseline** | | | |
|-------------------------|------------|----------|-------|--|
| (μ g/g dry wt.) | Tissue 🔵 | Sediment | | |
| Arsenic | 7.5 | 0 | 4.1 | |
| Cadmium | 1.90 | 0 | 1.00 | |
| Copper | 10 | | 31 | |
| Lead | 0.73 | | 22 | |
| Methyl Mercury | 0.01 | | 0.002 | |
| Mercury | 0.04 | | 0.105 | |
| Zinc | 54 | | 95 | |

| Organics | Baseline** | | | |
|-------------------|------------|----------|-------|--|
| (ng/g dry wt.)* | Tissue 🔵 | Sediment | | |
| Alpha-Chlordane | 8.5 | 0 | 0.58 | |
| Chlorpyrifos | 14 | • | 0.27 | |
| DDT | 29 | 0 | 3.0 | |
| Heptachlor | 0 | | 0.00 | |
| Hexachlorobenzene | 6 | 0 | 1.01 | |
| Mirex | 37.4 | | 1.060 | |
| Benzo[a]pyrene | 34 | 0 | 51 | |
| Benzo[e]pyrene | 288 | • | 45 | |
| PCB | 293 | | 18 | |

^{*}Tissue data is lipid normalized. **Core basin-wide measurements (AOC and reference sites).



Bervoets L., J. Voets, A. Covaci, S.G. Chu, D. Qadah, R. Smolders, P. Schepens, and R. Blust. 2005. Use of transplanted zebra mussels (*Dreissena polymorpha*) to assess the bioavailability of microcontaminants in Flemish surface waters. Environmental Science and Technology 39(6):1492-505.

Berny, P., O. Lachaux, T. Buronfosse, M. Mazallon, and G. Gillet, 2002, Zebra mussels (Dreissena polymorpha) as indicators of freshwater contamination with lindane. Environmental Research 90(2):142-51.

Bruner, K. A., S. W. Fisher, and P. F. Landrum. 1994a. The role of the zebra mussel, *Dreissena polymorpha*, in contaminant cycling .1. The effect of body-size and lipid-content on the bioconcentration of PCBs and PAHs. Journal of Great Lakes Research 20:725-734.

Bruner, K. A., S. W. Fisher, and P.F. Landrum. 1994b. The role of the zebra mussel, *Dreissena polymorpha*, in contaminant cycling. 2. Zebra mussel contaminant accumulation from algae and suspended particles. and transfer to the benthic invertebrate. Gammarus fasciatus, Journal of Great Lakes Research 20:735-750.

Carlton, J. 2008. The zebra mussel Dreissena polymorpha found in North America in 1986 and 1987. Journal of Great Lakes Research 34(4):770-773.

Cho, Y. C., R. C. Frohnhoefer, and G. Y. Rhee. 2004. Bioconcentration and redeposition of polychlorinated biphenyls by zebra mussels (Dreissena polymorpha) in the Hudson River. Water Research 38:769-777.

de Kock, W.C., and C.T. Bowmer. 1993. Bioaccumulation, biological effects and foodchain transfer of contaminants in the zebra mussel (Dreissena poymorpha). Pages 503-533 in Zebra Mussels: Biology, Impacts, and Controls, T.F. Nalepa and D.W. Schloesser, editors. Lewis Publishers. Baco Raton, FL.

Farrington, J.W., 1983. Bivalves as sentinels of coastal chemical pollution: the Mussel (and oyster) Watch. Oceanus 26(2):18-29.

GLRI Action Plan 2010. White House Council on Environmental Quality, "Great Lakes Restoration Initiative Action Plan for 2012-2014" (2010). Government Documents. Paper 1. http:// greatlakesrestoration.us/pdfs/glri actionplan.pdf

Great Lakes Water Quality Agreement 2012. Protocol amending the agreement between Canada and The United States of America on Great Lakes water quality, 1978, As Amended on October 16, 1983 and on November 18, 1987.

Gossiaux, D.C., P. F. Landrum, and S. W. Fisher. 1998. The assimilation of contaminants from suspended sediment and algae by the zebra mussel, Dreissena polymorpha. Chemosphere 36:3181-3197.

IJC 2011. Fifteenth biennial report of Great Lakes water quality. International Joint Commission, Canada and the United States. 59 pp. http://www.ijc.org/rel/boards/ watershed/15biennial report web-final.pdf

Kimbrough, K. L., and G. G. Lauenstein (eds.). 2006. Major and trace element analytical methods of the

National Status and Trends Program: 2000-2006. NOAA Technical Memorandum NOS NCCOS 29 Silver Spring, MD. 21 pp. http://ccma.nos.noaa.gov/publications/nsandtmethods.pdf

Kimbrough, K. L., G. G. Lauenstein and W. E. Johnson (eds.). 2006. Organic contaminant analytical methods of the National Status and Trends Program: Update 2000-2006. NOAA Technical Memorandum NOS NCCOS 30 Silver Spring, MD. 137 pp. http://www.ccma.nos.noaa.gov/publications/organicsmethods.pdf

Lauenstein, G. G. and A. Y. Cantillo. 1998. Analytical methods of the National Status and Trends Program Mussel Watch Project - 1993 -1997 Update. NOAA Technical Memorandum NOS ORCA 130. Silver Spring, MD. http://www.ccma.nos.noaa.gov/publications/tm130.pdf

MacDonald, D. D., C. G. Ingersoll, and T. A. Berger. 2000. Development and evaluation of consensus-based sediment quality guidelines for freshwater ecosystems. Archives of Environmental Contamination and Toxicology 39: 20-31.

Marvin C. H., E. T. Howell., and E. J. Reiner. 2000. Polychlorinated dioxins and furans in sediments at a site colonized by *Dreissena* in western Lake Ontario, Canada. Environmental Toxicology and Chemistry 19: 344-351.

Mills, E.L., R. M. Dermott, E. F. Roseman, D. Dustin, E. Mellina, D. B. Conn, and A. P. Spidle. 1993 Colonization, ecology, and population structure of the "quagga" mussel (Bivalvia: Dreissenidae) in the lower Great Lakes. Canadian Journal of Fisheries and Aquatic Sciences 50: 2305-2314.

Morrison, H.A., F. A. P. C. Gobas, R. Lazar, D. M. Whittle and G. D. Haffner. 1998. Projected changes to the trophodynamics of PCBs in the western Lake Erie ecosystem attributed to the presence of zebra mussels (*Dreissena polymorpha*). Environmental Science and Technology 32: 3862-3867.

Reeders, H. H., and A. Bij de Vaate. 1992. Bioprocessing of polluted suspended matter from the water column by the zebra mussel (*Dreissena polymorpha* Pallas). Hydrobiologia 239:53-63.

Robertson, A., and G. G. Lauenstein. 1998. Distribution of chlorinated organic contaminants in dreissenid mussels along the southern shores of the Great Lakes. Journal of Great Lakes Research 24(3):608-619.

Schantz, M.M., R.M. Parris, and S.A. Wise. 2008. NIST Intercomparison program for organic contaminants in the marine environment: Description and results of the 2007 organic intercomparison exercise. National Institute of Standards and Technology, Gaithersburg, MD. NISTIR 7501, pp279. Silver Spring, MD.

U.S. EPA. 2001. Environmental Monitoring and Assessment Program (EMAP): National Coastal Assessment Quality Assurance Project Plan 2001-2004. United States Environmental Protection Agency, Office of Research and Development, National Health and Environmental Effects Research Laboratory, Gulf Ecology Division, Gulf Breeze, FL. EPA/620/R-01/002.

Vanderploeg, H. A., T. F. Nalepa, D. J. Jude, E. L. Mills, K. T. Holeck, J. R. Liebig, I. A. Grigorovich, and H. Ojaveer. 2002. Dispersal and emerging ecological impacts of Ponto-Caspian species in the Laurentian Great Lakes. Canadian Journal of Fisheries and Aquatic Sciences 59: 209-1228. http://www.glerl.noaa.gov/pubs/fulltext/2002/20020012.pdf

Willie, S. 2007. NRC/20 Twentieth intercomparison for trace metals in marine sediments and biological tissues. National Research Council Canada. NRCC No. 50099. 72 pp.



U.S. Department of Commerce

Penny Pritzker, Secretary

National Oceanic and Atmospheric Administration

Kathryn Sullivan, Under Secretary for Oceans and Atmosphere

National Ocean Service

Holly Bamford, Assistant Administrator for Ocean Service and Coastal Zone Management









The mission of the National Centers for Coastal Ocean Science is to provide managers with scientific information and tools needed to balance society's environmental, social and economic goals. For more information, visit: http://www.coastalscience.noaa.gov/



